


*The Kentucky
Agricultural Experiment Station*

109th


Annual Report

1996

College of Agriculture
University of Kentucky • Lexington, Kentucky 40546




Contents

.....	7
Agricultural Forestry	9
Agroforestry	11
To His Excellency, The Honorable Paul Patton Governor of Kentucky	14
.....	16
.....	20
I herewith submit the one hundred and ninth annual report of the Kentucky Agricultural Experiment Station for the period ending December 31, 1996. This is done in accordance with an act of Congress, approved March 2, 1887, entitled, "An act to establish Agricultural Experiment Stations, in connection with the Agricultural Colleges established in the several states under the provisions of an act approved July 2, 1862, and under the acts supplementary thereto," and also the act of Kentucky State Legislature, approved February 20, 1888, accepting the provisions of the act of Congress.	23
.....	25
Very respectfully,	26
	28
C. Oran Little, Director	31
Lexington, Kentucky	34
June 30, 1997	35
Staff	36

To His Excellency
The Honorable Paul Patton
Governor of Kentucky

I have the honor to acknowledge the receipt of the report of the Kentucky Agricultural Experiment Station for the period ending December 31, 1906. This is done in accordance with an act of Congress, approved March 3, 1887, entitled "An act to establish Agricultural Experiment Stations in connection with the Agricultural Colleges established in the several States under the provisions of an act approved July 2, 1862, and under the act supplementary thereto," and also the act of Kentucky State Legislature, approved February 20, 1888, accepting the provisions of the act of Congress.

Very respectfully,

C. O. Curtis, Director
Washington, Kentucky
June 30, 1907

Contents

Purpose	7
Agricultural Economics	9
Agronomy	11
Animal Sciences	14
Biosystems and Agricultural Engineering	16
Entomology	18
Forestry	20
Horticulture and Landscape Architecture	22
Nutrition and Food Science	25
Plant Pathology	27
Regulatory Services	30
Robinson Substation	33
Rural Sociology	36
Veterinary Science	38
Publications	41
Statewide Research	54
Financial Statement	55
Staff	56

Contents

7	Purpose
9	Agricultural Economics
11	Agronomy
14	Animal Sciences
16	BioSystems and Agricultural Engineering
18	Entomology
20	Forest
22	Horticulture and Landscape Architecture
23	Nutrition and Food Science
25	Plant Pathology
30	Regulatory Services
33	Robinson Station
36	Rural Sociology
38	Veterinary Science
41	Publications
54	Statwide Research
55	Statwide Station
56	Stat

Agriculture Economics Purpose

As a Land Grant institution, the University of Kentucky is responsible for serving the people of the Commonwealth of Kentucky. The College of Agriculture, with its research, teaching, and extension activities, has developed a structure and organization to provide the mandated Land Grant services in agriculture and related areas.

The Kentucky Agricultural Experiment Station has been providing research results to farmers for more than 100 years. The continued growth of Kentucky agriculture attests to the benefits of applying new knowledge and technology to the agricultural production process. Much of the research leading to increased quantity and improved quality of Kentucky's agricultural output was performed by the Experiment Station. Also, College researchers address problems of agribusiness, consumers, international trade, food processing, nutrition, community development, soil and water resources, and the environment.

Although much Experiment Station research has immediate application to agricultural problems, scientists are also involved in basic research, generating new information to help solve present and potential problems. The ability of Kentucky producers to be competitive in domestic and world markets requires an expanded base of knowledge in emerging areas of research applicable to agriculture.

This Annual Report summarizes Experiment Station research highlights for 1996. Lists of the faculty, research projects, and publications completed during the year are also provided.

The research programs of the Kentucky Agricultural Experiment Station have benefited Kentucky's agriculture over the past century, and the results of present and future research will continue to serve Kentucky's primary industry.

Rural Economic Development Research - Telecommunications

Our TVA Rural Economic Development Studies Center is organized to support the development and dissemination of useful research for use by decision makers engaged in rural economic development. We have defined three main areas for our work. The Center's most visible product has been *The Farmer's Guide to the Internet*. The first edition of the book was printed in March 1996, and by the end of December more than 10,000 copies were sold. *The Farmer's Guide* has been purchased by individuals and small businesses in rural areas who seek a source of information on how to get access to the internet, and by large corporations who use it as a tool to use in internet training. In addition, the Center produces a research report with the National Rural Electric Cooperative Association on telecommunications opportunities for electric cooperatives. This study, "The New Telecommunications Environment," has been used in a series of regional workshops to help managers of rural electric cooperatives and municipal power systems assess opportunities for sup-

porting their telecommunications needs. The Center also provides technical assistance to rural electric cooperatives in the form of a "Rural Electric Cooperative Telecommunications Handbook." The handbook provides information on how to use telecommunications services to improve the efficiency of their operations. The handbook is available in both print and electronic formats. The Center also provides technical assistance to rural electric cooperatives in the form of a "Rural Electric Cooperative Telecommunications Handbook." The handbook provides information on how to use telecommunications services to improve the efficiency of their operations. The handbook is available in both print and electronic formats.

The Center's research on the impact of the 1997 heavy crop is also being completed. The Center's research on the impact of the 1997 heavy crop is also being completed. The Center's research on the impact of the 1997 heavy crop is also being completed.

Enhancing Farm Profitability

Analysis of more than 250 farms participating in the Kentucky Farm Business Analysis program revealed that 1993 was a record year. Financially, farmers varied around the state and by enterprise.

Purpose

As a Land Grant institution, the University of Kentucky is responsible for serving the people of the Commonwealth of Kentucky. The College of Agriculture, with its research, teaching, and extension activities, has developed a structure and organization to provide the mandated Land Grant services in agriculture and related areas.

The Kentucky Agricultural Experiment Station has been providing research results to farmers for more than 100 years. The continued growth of Kentucky agriculture attests to the benefits of applying new knowledge and technology to the agricultural production process. Much of the research leading to increased quantity and improved quality of Kentucky's agricultural output was performed by the Experiment Station. Also, College researchers address problems of agriculture, consumers, managers, and food processing, nutrition, community development, soil and water resources, and the environment.

Although much Experiment Station research has been applied to agricultural problems, wherever we also involved in basic research, generating new information to help solve present and potential problems. The ability of Kentucky producers to be competitive in domestic and world markets requires an expanded base of knowledge in emerging areas of research applicable to agriculture.

This Annual Report summarizes Experiment Station research highlights for 1996. Lists of the faculty, research projects, and publications completed during the year are also provided.

The research programs of the Kentucky Agricultural Experiment Station have benefited Kentucky's agriculture over the past century, and the results of present and future research will continue to serve Kentucky's primary industry.

Agricultural Economics

Research in Agricultural Economics involves diverse subjects including improving profitability, through management and marketing, environment and natural resources and assisting in the revitalization of rural communities.

Risk Management Research

The 1995/96 Farm Bill changed the way government will support farm prices and incomes for the next seven years, and quite possibly forever. Thus, when market prices go down, farmers cannot count on government payments to make up the difference. This will increase risk. Many argue that the changes will also increase price volatility. Recent crop failures also remind farmers that significant yield risk continues to be part of cash crop farming.

Crop insurance products are evolving to handle both price and yield risk. Work at the University of Kentucky in the past five years has been focused on new products for the crop insurance program. The Group Risk Plan (GRP) was developed at the University of Kentucky. The GRP is the only USDA program that attempts to adjust for technological advance by using trend county yields as the index to make payments to farmers. By insuring the combination of price and yield shortfalls, new products will provide more effective risk management for cash crop farmers. Economists at the University of Kentucky are involved in reviewing these products.

Rural Economic Development Research—Telecommunications

Our TVA Rural Economic Development Studies Center is organized to support the development and dissemination of useful research for use by decision makers engaged in rural economic development. We have defined three main areas for our work. The Center's most visible project has been *The Farmer's Guide to the Internet*. The first edition of the book was printed in March 1996, and by the end of December more than 10,000 copies were sold. *The Farmer's Guide* has been purchased by individuals and small businesses in rural areas who seek a source of information on how to gain access to the Internet, and by large corporations and universities to use in Internet training. In addition, the Center produces a research report with the National Rural Electric Cooperative Association on telecommunications opportunities for electric cooperatives. This study, "The New Telecommunications Environment," has been used in a series of regional workshops to help managers of rural electric cooperatives and municipal power systems assess opportunities for aug-

menting their electricity operations with telephone, cable or Internet subsidiaries.

Rural Economic Development Research—Environment

Experts around the world are currently debating how environmental conditions and regulations affect economic growth. Research recently completed at the University of Kentucky shows that states within the U.S. which had better environmental conditions also experienced more per capita income growth, all else being equal. This suggests that relationships between growth and the environment can be complementary, and that it is not necessarily always competitive as is commonly believed. This research has an impact on how researchers conceptualize and model the relationship between growth and the environment over time. The research also suggests that communities which strive to improve their natural environments will experience faster income growth than those which do not.

Tobacco Research

Agricultural Economics research into the cost-saving benefits of lower plant populations for burley tobacco is saving Kentucky tobacco farmers thousands of dollars in reduced labor costs. The collaborative extension educational efforts of agricultural economics, agronomy, agricultural engineering, and plant pathology specialists have resulted in some major shifts toward wider-spaced tobacco plants. Economic/agronomic research suggests that reducing per acre plant population by 25% can lower costs of production by eight cents per pound. Labor savings account for most of the cost advantage with per acre labor reductions in the 20- to 40-hour per acre range.

If half of Kentucky's 220,000-acre 1997 burley crop is grown at 6,500 plants per acre instead of 8,500 plants per acre, the labor savings alone could account for the equivalent of 1,320 full-time jobs. Or, from another perspective, an eight cents per pound cost reduction is worth \$22,000,000 in extra net farm income if even half of the 1997 crop is produced at wider spacings.

Enhancing Farm Profitability

Analysis of more than 280 farms participating in the Kentucky Farm Business Analysis program revealed that 1995 was a mixed year, financially. Returns varied around the state and by enterprise.

All Purchase area farms, Ohio Valley area livestock farms, Pennyroyal area grain and livestock farms, and Central Kentucky hog and crop farms improved upon returns in 1995, whereas Ohio Valley area grain farms, Pennyroyal area dairy farms, and Central Kentucky dairy and beef farms had lower returns than in 1995. The Pennyroyal livestock and dairy farms and the Central Kentucky dairy, hog, and beef farms experienced negative net returns. The operator's share of net farm income (showing the financial rewards for the operator, their equity capital, and their management input) was positive for all six groups in the state.

Livestock Marketing Research

A model of the livestock industry was developed that could track the reaction of beef cow numbers to changes in pastureland as well as dairy cow numbers, beef prices, and other variables internal to the industry. A quarterly version of this model helped to explain what response beef cattle prices would have to the dairy cow buyout of the mid-1980s. There were some pretty dire forecasts at that time, but our analysis suggested that the effect would be minimal. And so it proved.

The model was used to anticipate the effects of biotechnological innovations that are under development. Joint work with economists at the University of Georgia indicated that, with current feeder steer/heifer price spreads, cow-calf producers would find it profitable to adopt sexed technology to

determine the offspring gender under a wide range of cost estimates and treatment effectiveness rates.

Price effects would not wipe out profitability of adopting the technology at moderate adoption rates, up to 50%, and given our estimates of application costs. Steer prices do not decline as application rates rise, however. The analysis to date indicates that major price effects and changes in the structure of the beef cattle industry might result from widespread adoption of sexed-semen technology.

Research Projects

- Analyzing the Future International Competitiveness of the U.S. Food Industry — *M.R. Reed, M.A. Marchant, and L.L. Mather*
- An Analysis of Finance Efficiency and Socioeconomic Influences in Urban and Rural Public Schools — *S.J. Goetz and D.L. Debertin*
- An Evaluation of International Markets for Southern Commodities — *M.A. Marchant and M.R. Reed*
- Changing Patterns of Food Demand and Consumption Behavior — *B.W. Bobst*
- Development of Multiobjective Decision Criteria Models for Agricultural Investment Appraisal — *A. Pagoulatos and D.L. Debertin*
- Economic Analysis of Biotechnological Innovations in the U.S. Beef Cattle Industry — *B.W. Bobst*
- Economic and Environmental Impacts of Water-Quality Protection Policies on Kentucky Agriculture — *H.H. Hall and J.R. Skees*
- Evaluation of Public Policy Alternatives Designed to Help U.S. Cash Crop Farmers Manage Risk — *J.R. Skees and H.H. Hall*
- Regulatory, Efficiency and Management Issues Affecting Rural Financial Markets — *D. Freshwater*
- Rural Economic Development Alternatives in the New Competitive Environment — *S.J. Goetz and D. Freshwater*

Agronomy

The Department of Agronomy conducts research on soils, crop ecology/management, crop genetics, and plant biology. Our objectives include the improvement of plant quality and productivity for food, feed, and fiber applications. We strive to develop innovative plant production systems which are profitable, sustainable, and environmentally sound. To address these objectives, we require a diverse, interdisciplinary faculty with expertise ranging from cutting-edge molecular biology to applied production technology. The recent successes—and the tremendous future potential—of both crop biotechnology and precision agriculture systems make this a fast-moving and exciting time for agronomic science. Furthermore, the broad-based, popular concern for the quality of Kentucky's soil and water resources provides even greater urgency for research on sustainable land-management practices. Examples of significant research accomplishments during 1996 are described here.

- We are conducting several studies to exploit and evaluate the potential of the new precision agriculture/GPS technology. In one experiment, when corn population was varied according to depth of topsoil, returns per acre were increased relative to constant corn populations. This appears to be the first U.S. report of successfully controlling corn populations with a specific soil variable.
- We have introduced the animal gene for metallothionein, a compound which binds the heavy metal cadmium, into transgenic plants. We can target the gene to specific tissues in the plant. Expression of the gene was found to be most effective in the cytoplasm and perhaps in the vascular tissue of the plant.
- Our new wheat variety Foster will be available for production this year. Foster is a high yielding, high test weight variety with excellent disease resistance and superior milling and baking quality.
- A major research and education initiative on agronomic potential in eastern Kentucky is under way. Much of this work is sponsored by the Robinson Trust. A large beef cattle study on previously mined land will assess the impact of grazing on the plant community and soil fertility. A 27-acre site on the Laurel Fork of the Robinson Forest is being developed to demonstrate state-of-the-art reclamation technology.
- Weather-related losses during dry hay production and storage cost Kentucky livestock producers millions of dollars each year. We ensiled high-moisture forage in round bales using stretch-wrap plastic to achieve anaerobic conditions. In three years of studies, this increased nutritive value and reduced dry matter loss by 80% compared with hay stored outside.
- We continue to learn more about the mechanisms by which contaminants move through soil to springs and groundwater. We recently observed that soil colloids (very small particles of soil or organic matter) may be responsible for accelerating and increasing the leaching of heavy metals through the soil. In another study, we found that the supposedly inert surfactants applied with herbicides may have a strong influence on mobility and degradation of the herbicides.
- Soybean varieties with altered oils have been developed by private industry and are being grown in states to our north. Our work has shown that varieties with low linolenic acid level can be induced to produce even lower linolenic concentrations under our warmer Kentucky conditions. This suggests an opportunity for value-added soybean production in the near future.
- We have developed an efficient new regeneration system for soybean called somatic embryo cycling. It results in shorter cycling times for recovering plants and reduces detrimental effects from tissue culture. We believe this will be a major improvement for soybean transformation.
- Some of the first studies in the nation on the ability of forage grasses and alfalfa to withstand continuous, close grazing are in progress. These researchers are also cooperating with Plant Pathology and the Eden Shale Farm to study the effects of varietal resistance of alfalfa to aphanomyces root rot on alfalfa establishment.
- Two major advances in understanding plant-pathogen interactions were accomplished. The structure and function of a key phytoalexin enzyme has been analyzed in detail. This will advance rational design and genetic engineering of the genes coding for these enzymes. Further refinement in identifying the regulatory DNA sequences controlling expression of plant defense genes was also completed. Both of these studies could lead to major advances in engineering disease resistance for plants.
- Growing three soybean varieties, one from each maturity group II, IV, and V, provides greater yield stability than growing one superior variety or growing three varieties of the same maturity. Techniques were developed for selecting varieties that provide high yield and stability without changing varieties every year.
- We found that in late plantings, resistance to soybean mosaic virus provided a 12% yield advantage compared to SMV-susceptible soybean. For double-cropped soybeans grown after winter wheat in Kentucky, soybean producers should benefit by selecting high yielding varieties with SMV resistance.
- We have generated a large number of transgenic tobacco lines with different pathogen-derived resistance genes.

Tobacco lines resistant to tobacco vein mottling, tobacco etch, potato virus Y, alfalfa mosaic, and tomato spotted wilt viruses have been produced. These germplasm are being greenhouse and field evaluated for virus resistance and agronomic traits prior to deployment in future cultivars.

- In collaboration with the Kentucky Tobacco and Health Research Institute, several Agronomy faculty are pursuing possibilities for extended uses of tobacco. It has been demonstrated that genetically engineered tobacco can produce a variety of valuable biochemicals including antibodies, vaccines, enzymes, and industrial feedstocks. Further research in molecular biology, product engineering, and economic analysis will be required to bring these opportunities to the farm.
- Our turfgrass project on fungal disease control via Integrated Pest Management has been investigating the effects of dewfall on golf course greens and fairways. We measured leaf surface moisture to determine the amount of accumulation and re-accumulation after removal on creeping bentgrass. This knowledge of how and when moisture accumulates on turfgrasses will help managers reduce the duration of leaf wetness, thus reducing disease pressure without the use of fungicides.

Research Projects

A Silica/Fly Ash Based Technology for Controlling Pyrite Oxidation — *V.P. Evangelou*
 Agri Industry Development Program — *C.W. Absher and M. Rasnake*
 Alkaloid Accumulation in *Acremonium Coenophialum* Infected Tall Fescue — *L.P. Bush and H.R. Burton*
 Altered Fatty Acid Varieties: Does Our Climate Give Us an Angle? — *L.J. Grabau*
 Altering Ergot Alkaloid Biosynthesis by the *Acremonium* Endophyte of Tall Fescue — *C.L. Schardl, M.R. Siegel, and L.P. Bush*
 Ameliorative Designs to Improve the Efficiency of Constructed Wetlands Treating High Metal Load Acid Mine Drainage in the Rock Creek Watershed — *A.D. Karathanasis*
 Analysis of MRNA Polyadenylation and Metabolism in Plants — *A.G. Hunt*
 Backup of Clover (CLO) Accessions in the NSSL — *N.L. Taylor*
 Backup of Clover (CLO) Accessions in NSSL - Phase 2 — *N.L. Taylor*
 Bacterial Dissimilation of Nitrate to Ammonium in Batch and Chemostat Culture — *M.S. Coyne*
 Behavior, Fate, and Bioactivity of Acetolactate (ALS)-Inhibiting Herbicides — *M. Barrett and W.W. Witt*
 Biochemistry and Molecular Biology of Sesquiterpene Cyclase and Squalene Synthetase from Tobacco — *J. Chappell*
 Breeding Burley Tobacco for Improved Pest Resistance and Productivity — *M.T. Nielsen, B.S. Kennedy, and P.D. Legg*
 Breeding Improved Varieties of Wheat, Oats, and Barley for Kentucky — *D.A. Van Sanford*
 Cellular and Molecular Biology Initiative in Dark Tobacco — *G.B. Collins*
 Characterization and Classification of Kentucky Soils — *A.D. Karathanasis and R.I. Barnhisel*
 Characterization and Modification of Heavy Metal Accumulation in Plants, with Emphasis on Tobacco — *G.J. Wagner*
 Characterization of Phytoalexin and Sterol Biosynthetic Genes in Tobacco — *J. Chappell*
 Characterization of a Plant Poly A Polymerase — *A.G. Hunt*
 Classifying Soils for Solute Transport as Affected by Soil Properties and Landscape Position — *E. Perfect*
 Cloning Epoxy Fatty Acid Genes — *D. Hildebrand*

Cloning and Heterologous Expression of Pesticide Metabolizing Cytochrome P450 Genes — *M. Barrett*
 Cloning Genes Encoding Enzymes for Epoxy Fatty Acid Accumulation in Oilseeds — *D. Hildebrand*
 Corn Breeding and Genetics: White Endosperm Breeding, Food Quality Inheritance, and Hybrid Performance Tests — *C.G. Poneleit*
 Correlation and Calibration of Crop Yields with Soil Test Levels of Major Nutrients — *W.O. Thom*
 Cropping and Planting Systems to Allow Economic Canola Production — *J.H. Herbek and L.W. Murdock, Jr.*
 Damage Thresholds Risk Assessment and Environmentally Compatible Management Tactics for White Grub Pests of Turfgrass — *D.A. Potter, K.F. Haynes and A.J. Powell, Jr.*
 Dark Tobacco Breeding Genetics and Management — *P.D. Legg*
 Determining Optimum Rates for Several Crop Nutrients — *W.O. Thom*
 Development of a Basic Soil Morphology Training Course for Onsite Sewage Disposal Treatment System Personnel — *A.D. Karathanasis*
 Development of Efficient Tissue Culture and Genetic Engineering of Soybean — *G.B. Collins and R. Dinkins*
 Direct Vegetation of Fly Ash — *R.I. Barnhisel*
 Distribution of Constituents within Tobacco Leaf — *H.R. Burton and L.P. Bush*
 Does Kentucky Need an Early Maturing Soybean Variety Trial? — *L.J. Grabau*
 Early Maturing Soybean Cropping System: Identifying Appropriate Cultivars — *L.J. Grabau*
 Effect of Row Width on Corn Yields — *M.J. Bitzer and J.H. Herbek*
 Effects of Riparian Vegetation on Water Quality: Modeling and Experimental Studies — *B.J. Barfield, R.L. Blevins, V.P. Evangelou, and D.T. Carey*
 Effects of Suckering Practices on Growth Characteristics — *J. Calvert*
 Environmental and Genotypic Control of Assimilate Allocation in Grain Crops — *D.B. Egli*
 Environmental and Morphological Determinants of Field Curing Rates of Legume Hay — *M. Collins and N.L. Taylor*
 EPA/EPSCOR: Assessing Transport of Colloid Bound Herbicides and Heavy Metals to Groundwater — *A.D. Karathanasis, R.E. Phillips*
 Establishing an Advanced Techniques Course in Biotechnology — *J. Chappell*
 Evaluation of Burley Tobacco Varieties — *M.T. Nielsen, J. Calvert, and B. Kennedy*
 Evaluation of Soybean Varieties and Breeding Lines for Use in Kentucky — *T.W. Pfeiffer, J.M. Wood, and C.R. Tutt*
 Field Application of Pyrite Microencapsulation Technologies for Controlling Pyrite Oxidation and Acid Mine Drainage Production — *V.P. Evangelou and R.E. Phillips*
 Forage Crop Breeding to Improve Yield and Quality — *N.L. Taylor*
 Foreign Gene Introduction into Soybean — *G.B. Collins and R. Dinkins*
 Fungal Pathogen Resistance in Dark Tobacco — *M.T. Nielsen*
 Genetic Control of a Putative Multiple Pesticide Metabolizing Cytochrome P-450 — *M. Barrett*
 Genetic Engineering of Dark Tobaccos: A Sub-project of Cellular and Molecular Biology Initiative in Dark Tobacco — *J. Chappell*
 Genetic Engineering of Soybeans for Increased Oil Content and Epoxy Fatty Acid Accumulation — *D. Hildebrand*
 Genetic Engineering of Soybeans for Increased Value — *D. Hildebrand*
 Genetic Improvement of Soybean for Kentucky — *T. Pfeiffer*
 Grain Quality Laboratory — *C.G. Poneleit*
 HHMI Initiative Teacher Research — *T.D. Phillips*
 Identification Characterization Seed Increase of Clovers — *N.L. Taylor*
 Improved Fertilizer Use Efficiency and Environmental Soundness in Burley Tobacco Production — *J.L. Sims*
 Improved Management of Legume Cover Crops for Sustainable Grain Crop Production — *W.W. Frye, R.L. Blevins, and J.H. Herbek*
 Improving Switchgrass Productivity as a Biofuel Crop — *M. Rasnake*
 Increased Desaturation of Soybean Triacylglycerols — *D. Hildebrand*
 Influence of Soil Structural Heterogeneity on Transport of Fecal Pathogens and Solutes with the Vadose Zone — *E. Perfect*
 Isolation of Sclareol/Labdenediol Synthase — *G.J. Wagner*
 Kinetics and Thermodynamics of Adsorption-Desorption in Binary and Ternary Soil Colloid Systems — *V.P. Evangelou*

- Maintenance of Seed of Trifolium Species — *N.L. Taylor*
 Mechanisms for Vacuolar Storage/Sequestration of Cd, Zn, Mn, Ni — *G.J. Wagner*
 MOA #13814 with Natural Resources and Environmental Protection Cabinet Division of Water — *A.D. Karathanasis*
 Modifying Recombination Rates in Soybean and Assessing the Effect on Breeding Progress — *T.W. Pfeiffer*
 Multiplicative (Linear-Bilinear) Models for Genotype X Environment Interaction in Crop Cultivar — *P.L. Cornelius*
 Multiplicative Models for Genotype X Environment Interaction — *P.L. Cornelius*
 Nicotine Synthase and Nicotine Demethylase — *L.P. Bush*
 On-Farm No-Till Studies: Making It Work — *L.J. Grabau and J.H. Grove*
 Pest Control Strategies for Grazing Livestock using Grass Endophyte Associations — *C.T. Dougherty, F.W. Knapp and L.P. Bush*
 Phenology, Population Dynamics, and Interference: A Basis for Understanding Weed Biology and Ecology — *W.W. Witt*
 Plant, Animal, and Environmental Factors Limiting Intake of Grazing Beef Cattle — *C.T. Dougherty*
 Plant Genetic Resource Conservation and Utilization — *N.L. Taylor*
 Polishing Peers: Improving Plant and Soil Science Education through Peer Review — *L.J. Grabau, M.S. Smith, L.K. Worley, and W. Burke*
 Population Improvement and Line Development of White Endosperm Maize — *C.G. Poneleit*
 Potyvirus Replication and Pathogenicity — *A.G. Hunt*
 Predicting Changes in Corn Seed Quality during Storage — *D.M. TeKrony*
 Predicting Nitrogen Needs of Wheat by Measuring Chlorophyll — *L.W. Murdock, Jr.*
 Refining Components of an Early-planted, Early-maturing Soybean Cropping System — *L.J. Grabau*
 Relationship between Photosynthesis, Assimilate Supply, and the Size of the Reproductive Sink — *D.B. Egli*
 Restoration of Altered Lands — *R.I. Barnhisel*
 Screening Soft Red Winter Wheat Varieties and Breeding Lines for Resistance to Head Scab — *D.A. Van Sanford*
 Seed Biology and Technology Investigations — *D.M. TeKrony and D.B. Egli*
 Soft Red Winter Wheat Breeding and Variety Development for Kentucky — *D.A. Van Sanford*
 Soil and Crop Nitrogen Testing to Improve Nitrogen Management for Burley Tobacco — *R.C. Pearce, J.G. Grove, and D.C. Ditsch*
 Soil Nitrate Testing to Improve N Use Efficiency and Reduce Residual Nitrate under Corn and Wheat — *J.H. Grove, D.A. Van Sanford, and C.G. Poneleit*
 Soil Survey Characterization and Environmental Impact Assessment of Daniel Boone National Forest Ecosystems — *A.D. Karathanasis*
 Somatic Cell Genetics of Crop Plants — *G.B. Collins*
 Soybean Tissue Culture and Genetic Engineering Center — *G.B. Collins*
 Studies of a Novel Pathway for the Biosynthesis of Straight and Branched, Odd and Even Length, Medium-chain Fatty Acids in Plants — *G.J. Wagner*
 Studies to Reduce Cadmium Accumulation in Tobacco and Research toward Extending the Uses of Tobacco — *G.J. Wagner*
 Targeting of the Cd-Chelator Metallothionein to the Plant Cell Wall and Root Tissue Using Recombinant DNA Methodology — *G.J. Wagner*
 Terpene Cyclases: Functional Domains and Structures — *J. Chappell*
 319 Program Assessment of Constructed Wetlands for Animal Waste Phase II — *W.O. Thom, Y. Wang, and J. Dinger*
 319 Program Implementation and Demonstration of BMPs for the Utilization of Poultry Litter in the Lower Green River Water Shed — *C.W. Absher, G. Henson, and W.O. Thom*
 319 Program Site-specific Nutrient and Biosolids Management on Agricultural Lands — *R.I. Barnhisel and S.A. Shearer*
 Understanding and Manipulation of Lipid Biosynthesis in Plants — *D.F. Hildebrand*
 Understanding Recombination and Modifying its Frequency in Soybean and Corn — *T.W. Pfeiffer and C.G. Poneleit*
 Using Early-maturing Soybean Varieties to Help Manage Soybean Cyst Nematode — *L.J. Grabau*
 Using Farm Family Case Studies to Teach Sustainable Agriculture — *M. Rasnake*
 Utilization of Coal Combustion By-products in Agriculture and Reclamation — *W.O. Thom*
 Utilizing Seed Vigor as a Component of Seed Quality — *D.M. TeKrony*
 Variable Rate Seeding: A Cooperative Study with Kentucky Corn Producers — *R.I. Barnhisel, S.A. Shearer, and M.J. Bitzer*
 Yield Evaluation of Alfalfa Varieties — *L. Lauriault and J.C. Henning*
 Yield Potential and Long-term Effects of No Tillage on Wheat Production — *L.W. Murdock, Jr., J.H. Herbek, and J.R. Martin*

Animal Sciences

The Department of Animal Sciences conducts research on poultry, swine, sheep, dairy cattle, beef cattle, horses, and foods. The research enhances our teaching program by providing new, up-to-date information for instructors and educational experiences for students. We conduct research from the molecular level to livestock production systems. The discovery research is designed to elucidate chemical and physiological functions that will eventually improve efficiency of livestock performance and enhance the quality and safety of our food supply.

The Cooperative Extension Service disseminates research results that can be applied to livestock production, food handling, and food processing. We are fortunate to have a system that allows teaching, research, and extension personnel to work together to provide new knowledge to keep us self-sufficient in our food supply. Many countries with dense population, such as China, will never be self-sufficient in food production since an extremely low percentage of their land is tillable and much of that is close to highly urbanized areas.

Food production will become more globalized. Technological advances will be needed to feed the population. Research is the key to maintaining an adequate supply of safe, high-quality food, and livestock production is a necessity. The Department of Animal Sciences conducts research in several disciplines to help achieve the goal of providing an adequate supply of food. We strive to provide the technology for livestock producers and food processors in the Commonwealth of Kentucky to produce and process that food in a sustainable manner.

Food Science

Weakening of gels made from salt-extracted muscle proteins was not diminished by protease inhibitors, suggesting that proteases were not involved in binding of meat particles during processing and cooking. Beef heart surimi prepared in the presence of the antioxidant propyl gallate and then blended with cryoprotectants had excellent long-term frozen storage stability and functionality. Antioxidant treatments retarded lipid oxidation during storage of both restructured and intact steaks, and a beefy flavoring masked the grassy off-flavor characteristics of grass-fed beef.

Studies were conducted to determine the effects of natural volatile compounds on bacteria *in vitro* and on the microflora of strawberry and blackberry fruits during storage. Percent reductions in populations of the eight test species varied with compound and exposure and ranged from 0% to 100% after 2, 4, and 6 h exposure. Microbial populations of straw-

berry and blackberry fruits decreased during storage at 2°C and increased after transfer to 22°C. Natural volatile compounds showed antimicrobial properties which may be useful in controlling microbial growth during storage of some foods.

Research was conducted on flavor problems associated with soy protein products (the single greatest impediment to their use in human foods). As part of this research, we discovered a major contributor to the undesirable flavor of soy protein. The mean content of 2-pentyl pyridine was found to be 0.65 ppm which is 54,000 times above its flavor threshold (0.000012 ppm). This strong flavor impact combined with its extremely repulsive flavor profile makes 2-pentyl pyridine a major contributor to the undesirable flavor of SPI.

Rumen Microbiology

Experimental models have been developed which explain the beneficial effects of several microbial supplements and competitive exclusion products on microbial populations in the gastrointestinal tract of domestic livestock. These models are based on the stimulation of critical physiology groups of bacteria and the elimination of specific pathogenic bacteria in the digestive tract. One of these models suggests that stimulatory effects of some of the yeast strains used in dietary supplements can be explained by the presence of several low molecular weight compounds which serve as growth modulators for anaerobic bacteria. Similar stimulatory compounds have not been previously reported.

Most ruminal microorganisms derive energy via the fermentation of carbohydrates. Our studies with the important ruminal bacteria *Ruminococcus albus* and *Prevotella ruminicola* indicated that both phosphorylytic and hydrolytic enzymes are used to metabolize the products of starch and cellulose degradation. In addition, we demonstrated that *P. Ruminicola* converts up to 40% of maltose into intracellular glycogen using a unique enzymatic pathway. This large deposition of glycogen was not previously recognized, and since *P. Ruminicola* is a predominant rumen organism this observation has potential implications for our understanding of ruminant nutrition.

Physiology and Breeding

The impact of heat stress or parturition on the antioxidant capacity of plasma from dairy cattle was evaluated. Heat stress depressed the antioxidant activity of plasma, and recovery of activity occurred when temperatures returned to thermoneutral levels. Antioxidant activity was also depressed

at calving, coincident with decreased vitamin E levels in plasma. Decreased antioxidant activity may be related to compromised resistance and increased incidence of mastitis during periods of heat stress in the periparturient period.

Although problems are associated with utilization of Brahman bulls, our regional results indicated no distinct overall advantage for alternative subtropically adapted sire breeds over the Brahman. Most of the alternative subtropically adapted sire breeds are expected to create less calving difficulty due to their siring calves with lighter birth weights; however, these calves are also expected to weigh less at weaning.

Swine Nutrition

Nitrogen excretion was reduced and ammonia concentration in anaerobic pits was suppressed when pigs were fed low-protein diets supplemented with amino acids. The immunoglobulin fraction of spray-dried porcine or bovine plasma increased growth performance of pigs weaned at 2 or 3 weeks of age, and pigs given exogenous somatotropin required higher dietary levels of a near-ideal blend of amino acids to maximize protein deposition in the carcass.

New information has been generated on the bioavailability of phosphorus in feedstuffs commonly fed to pigs. This new information allows nutritionists to more precisely formulate diets to meet pigs' available phosphorus requirements. Feeding low-phosphorus diets with added phytase improved bone mineralization and significantly reduced the excretion of phosphorus in the manure. Deletion of inorganic phosphorus from the diet of finishing pigs had little negative effect on performance or bone breaking strength, but it markedly reduced phosphorus excretion.

Ruminant Nutrition

Daily feed intake of lactating ewes was reduced for 7 days prior to weaning, and 96 mg of methscopolamine (MB) was injected at weaning. Both procedures effectively reduced milk production. Even greater reduction was achieved when MB administration was combined with feed restriction.

Data were collected on 276 cow-calf pairs grazing endophyte-infected tall fescue and supplemented with a molasses-based supplement continuously throughout each year. Supplemented cows produced heavier calves at weaning while maintaining heavier weights and higher body conditions.

Polypay wethers were allowed ad libitum access or were limit-fed soybean meal or molasses-supplemented diets. Based on gain, feed intake, and carcass data, molasses-based

rumen bypass supplement can replace equivalent soybean meal in growing-finishing diets without decreasing lamb performance or carcass characteristics.

Steers were used to determine the effect of small intestinal adaptation to a partial beta-amylase starch hydrolysate on activity and location of small intestinal glucose transport. Sodium-glucose cotransport activity was not affected by treatment, but did change along the small intestine. Maltase activity was maintained throughout the small intestine, whereas sodium-glucose cotransport activity may limit small intestinal starch assimilation in the distal small intestine. It does not appear glucose arising from carbohydrate hydrolysis regulates activity of sodium-dependent glucose transport in cattle.

Research Projects

- Acid-base Balance and Mineral Requirements of Dairy Cattle — *R.W. Hemken*
- Amino Acids Sources and Concentrations for Swine — *G.L. Cromwell*
- Beef Cattle Grazing: Endophyte-infected Tall Fescue with Alfalfa and Water Quality in Stream Pastures — *B.T. Larson*
- Bioenergetics of Nutrient Transport and Growth of Gram-negative Ruminant Microorganisms — *H.J. Strobel*
- Comparison of Forage Finishing Systems, Carcass Traits, and Processing Technologies — *W.G. Moody*
- Effect of Additives and Processing Methods on Culture Agglutination and Cheese Yield — *C.L. Hicks*
- Endocrine Mechanisms Contributing to Establishment of Pregnancy in Ruminants — *W.J. Silvia*
- Enhancing Food Safety through Control of Foodborne Disease Agents — *B.E. Langlois*
- Evaluation of Tall Fescue in Dairy Cattle — *J.A. Jackson*
- Functional Properties of Food Proteins — *Y.L. Xiong*
- Genetic and Phenotypic Aspects of Cow Productivity Using Field Records Collected on Angus Cattle — *D.K. Aaron*
- Hydrolyzed Feather Meal as a Supplement for Lambs Consuming High Concentrate on High Roughage Diets — *D.G. Ely*
- Impact of Level of Prewearing Performance on Subsequent Cow Herd Reproduction — *F.A. Thrift*
- Increased Efficiency in Sheep Production — *D.G. Ely*
- Induction of Puberty Onset in Beef Cattle — *K.K. Schillo*
- Lipid-derived Flavors/Odors and Their Association with Food Proteins — *W.L. Boatright*
- Metabolic Relationships in Supply of Nutrients for Lactating Cows — *D.L. Harmon*
- Microbial Strategies for Improving the Efficiency of Ruminant Production by Enhancing Propionate Metabolism in the Ruminant — *K.A. Dawson*
- Nutrition and Exercise on Development of Horse Skeletal and Muscular Tissue and Subsequent Performance — *L. Lawrence*
- Nutritional Systems for Swine to Increase Reproductive Performance — *M.D. Lindemann*
- Optimizing Digestion and Absorption in the Ruminant Small Intestine — *D.L. Harmon*
- Requirements and Bioavailability of Phosphorus for Swine — *G.L. Cromwell*
- Resistance to Mastitis in Dairy Cattle — *R.J. Harmon*
- Skeletal Problems in Poultry — *A.H. Cantor*

Biosystems and Agricultural Engineering

Biosystems and Agricultural Engineering research is directed toward solving existing and emerging engineering-related problems found on Kentucky's farms and forests, as well as developing methods of protecting foods and other farm products which are consumed or used by the public. Five broad areas of interest are pursued.

Machine Systems design involves development and evaluation of basic machine systems through the application of theoretical and applied mechanics. A fully-automated burley tobacco harvesting and curing system has been developed and is being tested on the Experiment Station. A mechanical burley tobacco spearing machine has been developed and demonstrated to farmers. A commercial version of the wire-strung portable frame system was used on a limited number of farms throughout the burley producing area. A two-row mechanical tobacco topper was shown to significantly reduce labor. An electric power tobacco stripping wheel has been developed that saves labor at the rate of three to five cents per pound.

Techniques for targeted herbicide application using GIS/GPS and reflectance sensors are being researched. A significant effort is in the area of robotics and machine vision as a support technology for machine systems for harvesting, grading, and automated control of field machinery. The effects of soil compaction created by heavy machinery on water infiltration, ground water movement, and plant growth are being studied.

Bioenvironmental Engineering is the application of principles of mathematics, chemistry, biology, and physics to sustain and improve the quality of our natural resources. The broad goal of current research efforts is to develop technology that improves the compatibility between water/soil resources and activities such as agriculture and mining. Research to minimize erosion and stream quality impacts from mined lands continues to be a productive, highly visible program involving improved detention basin design and continued development of computer-based design aids. Recently initiated research is devoted to assessing and reducing the effects of cattle grazing and tobacco production on runoff and subsurface quality. Increasing concerns regarding the environmental impacts of lawn care have prompted a study to relate runoff of commonly used herbicides to factors such as application rate, post-application irrigation, and storm severity.

Bioprocess Engineering involves optimization of equipment, sensors, and control algorithms for processes which use living cells or subcomponents of cells as bioreactors or biocatalysts. The cells typically employed include microbes, plant tissue cells, or mammalian cells and may be modified through biotechnology. Current research in this area includes optimizing the fermentation of *Aspergillus* to produce extra-

cellular enzymes for use as additives in animal feeds. The enzymes produced include a cellulase and hemicellulase to aid in cellulose digestion, and phytase to increase the efficiency of inorganic phosphorus use and reduce the phosphorus excreted by the animal. Another current research project is investigating the use of whole cells in organic solvents to catalyze reactions.

Structures and Environment involves the design of farm structures and environmental control systems for plant and animal production, feed storage and processing centers, residences, and utility buildings. Research efforts are directed toward reducing infiltration into residences, better management of animal waste, improved design of grain storage systems and structures, improved environmental control within poultry and swine growing facilities, computer-aided design methods for dairy facility design, and the development of alternative structures for curing burley and dark tobacco. Evaluation of a portable frame and field curing system for burley tobacco showed significant labor savings and excellent potential for adoption.

Crop Processing research involves basic engineering sciences, particularly heat and moisture transfer processes, to the processing, storage, and handling of farm products. Management protocol has been developed for curing burley tobacco in the field under plastic in various types of frameworks.

Food Engineering involves applying principles to achieve efficient production and high standards of quality during processing, packaging, storage, and distribution of food products. A milk coagulation sensor has been developed using fiber optics and light reflectance that improves the control of cheese making. The sensor is being evaluated in this country and several foreign countries. Experiments have been conducted that show the applicability of using light reflectance as an aid in making cottage cheese.

Other ongoing research which has basic implications in more than one of the areas mentioned above includes developing a profitable beef-forage production system through computer modeling and modeling growing swine. Meteorological research will improve the understanding and use of weather-related agricultural management models in the southern region. The Agricultural Weather Center provides: 1) weekly Kentucky weather summary for the National Weather Service for the Weekly Divisional Averages (WDA) for the Palmer Drought Index Model and dissemination on the National Weather Wire System and 2) Kentucky rainfall, temperature, and deviation-from-normal maps created daily for various time periods using the Geographical Information System (GIS) and made available on the World Wide Web. Research is under way to identify ways of reducing the health and accident risk for farm workers and youth.

Research Projects

- A System for Reconstruction or Rehabilitation of Rooting Media for Growing High-value Trees — *L.G. Wells*
- A Systematic Approach to Enzyme Recovery from Solid-state Fermentation — *S.E. Nokes*
- Agricultural Pesticide Handling and Application Technology Demonstrations — *S.G. McNeill*
- Analysis and Management of Misting Systems for Tunnel-ventilated Broiler Housing — *R.S. Gates*
- Assessment of the Hydrologic Response of Reclaimed Surface-mined Lands in the Appalachian Coal Region — *R.C. Warner*
- BMP Equine Waste Demonstration Project — *R.C. Warner*
- Comparison of Integrated Electronic Controllers to Conventional Staged Thermostatic Control in Large Broiler Houses — *R.S. Gates*
- Design and Demonstration of a Low-pressure Low-flow Irrigation System for Nurseries' Containerized and Ball and Burlap Areas — *R.C. Warner*
- Determination of Design Weather Parameters International Locations — *D.G. Colliver*
- Determination of Infiltration Characteristics of Mine Spoil Planted with Hardwoods — *R.C. Warner*
- Determination of Straw Properties and Air Flow Through an Upflow Biomass Gasifier — *W.E. Murphy*
- Determination of the Hydrologic and Sedimentation Response of Loose and Compacted Mine Spoil Planted with Hardwoods — *R.C. Warner*
- Development and Evaluation of a Model for an Active Gas Collection System at Municipal Landfills — *R.C. Warner*
- Development of a 3-axis Manipulator for Harvesting Fruit and Vegetable Crops — *S.A. Shearer*
- Development of Analytical Methods to Predict Stream/Aquifer Interactions in Alluvial Valleys — *S.R. Workman*
- Development of a Trickle Irrigation Design and Specification Program — *R.C. Warner*
- Development of Geographical Information Systems (GIS) in the UK College of Agriculture — *K.T. Priddy*
- Development of Near-real Time Weather Station at Woodford County Research Farm — *K.T. Priddy*
- Development of Profitable Beef-forage Systems for the Southern Region — *L.W. Turner*
- Development of Weather-related Insect Forecast Model Output in Geographical Information System (GIS) Format — *K.T. Priddy*
- Dietary Manipulation to Reduce Aerial and Effluent Nitrogen Concentrations in Broilers — *R.S. Gates*
- Dynamic Mist Control Strategies for Poinsettia Propagation — *R.S. Gates*
- Dynamic, Probabilistic Modeling of Respiratory Disease in Livestock — *L.W. Turner*
- Economic and Environmental Impacts of Water Quality Protection Policies on Kentucky Agriculture — *J.L. Taraba*
- Electric 4-H Project Manuals — *G.A. Duncan*
- Energy Savings from Combined Natural/Forced Ventilation Systems for Greenhouse Retrofits — *R.S. Gates*
- Enhanced Design of Burley Sparring Machine — *G.A. Duncan*
- Eroded Particle Size Distribution for Forest Soils, Mine Spoils, and Weathered Mine Spoils — *R.C. Warner*
- Estimating the Impact of Commercial Building Occupancy and Hot Water Needs in Sizing Ground Source Heat Exchanger Fields — *W.E. Murphy*
- Evaluation of a Bag-type Geothermal Ground Heat Exchanger — *W.E. Murphy*
- Facility Design and Testing for Closed System Plant Micropropagation — *R.S. Gates*
- Field Test of Low-cost Greenhouse Microprocessor Controls for Bedding Plant and Tobacco Plug Production — *R.S. Gates*
- GPS Tracking of Animal Position for Improved Pasture Systems Design/Layout — *L.W. Turner*
- Improved Information Delivery by Developing Internet Gopher/Web Server in the UK College of Agriculture — *K.T. Priddy*
- Improved Quality and Efficiency of Burley Tobacco Market Preparation — *L.G. Wells*
- Influence of Construction Quality Control on the Permeability of Soil Liner Waste Lagoons — *R.C. Warner*
- Interior Environment and Energy Use in Poultry and Livestock Facilities — *R.S. Gates*
- Inventory of Constructed Wetlands for Residential Treatment Systems — *R.C. Warner*
- Knowledge-based System for Single-stem Greenhouse Rose Production — *R.S. Gates*
- Landfill as a Bioreactor — *R.C. Warner*
- Measurement of Factors that Influence Soil Development, Water Transport, and Tree Growth in Reclaimed Surface Mines in Eastern Kentucky — *S.R. Workman*
- Mechanics of Granular Solids — *I.J. Ross*
- Milk Coagulation Sensor Development — *F.A. Payne*
- Model Development and Calibration for Predicting the Persistence and Efficacy of Chlorothalonil on Tomatoes — *S.E. Nokes*
- Model Development and Verification of a Tubular Trickle Irrigation System to Increase Water Application Efficiency — *R.C. Warner*
- Modeling of Heat Transfer from U-tube Heat Exchangers for Ground-source Heat Pumps — *W.E. Murphy*
- Modeling Responses of Growing Pigs — *L.W. Turner*
- Monitoring Impacts of Animal Research Center on Surface and Groundwater Quality — *J.L. Taraba*
- Nighttime Ventilation Strategies for Summer Heat Stress Relief in Broilers — *R.S. Gates*
- Optimization of Process Variables for Solid-state Fermentation Production of Xylanase — *S.E. Nokes*
- Physical and Bioremediation of Hazardous Waste Contained Soils through Incorporation of Coal Fines and In Situ Environmental Management — *R.C. Warner*
- Potential Utilization of Clay Deposits Extracted During Mining for Fly Ash Monofills — *R.C. Warner*
- Reducing Heat Stress in Dairy Cows through Forced Evaporative Cooling — *L.W. Turner*
- Reducing Nitrogen, Phosphorus, and Ammonia Erosion in Swine Waste through Diet Manipulation — *L.W. Turner*
- Reflectance Profile in Milk as a Function of Fat Content — *F.A. Payne*
- Residential Air Infiltration and Air Quality — *D.G. Colliver*
- Response of Hardwoods Planted in Mine Spoil and Receiving Trickle Irrigation — *R.C. Warner*
- Sensor Development for Cottage Cheese and Yogurt Culture — *F.A. Payne*
- Sequences of Extreme Temperature and Humidity for Design Conditions — *D.G. Colliver*
- Simulation of Infiltration and Lateral Subsurface Transport of Water in Hillslopes on Small Watersheds in Kentucky — *S.R. Workman*
- Site-specific Management of Herbicides on Crop Lands — *S.A. Shearer*
- Site-specific Crop Management: Nutrient Management — *S.A. Shearer*
- Stabilization of Embankments on AML Slopes Using Soil Bioengineering Techniques: A Field Evaluation of Cost-effectiveness — *R.C. Warner*
- Sterilizable Fiber Optic Probe for a Septic Food Processing Application — *F.A. Payne*
- Stress-strain Modeling of Wheel Compaction Incurred during Restoration of Prime Farmland — *L.G. Wells*
- Subsurface Leaching Potential of Animal Waste Holding Ponds as a Function of Soil Moisture and Compaction — *R.C. Warner*
- Testing of Time-integrated Variable Control in Tunnel-ventilated Broiler Housing — *D.G. Overhults*
- The Impact of Agricultural Systems on Surface and Ground Water Quality — *D.R. Edwards*
- Time-integrated Variable Control Strategies for Animal and Plant Environments — *R.S. Gates*
- Utilization of Commercially Composted Waste to Reduce the Transport of Herbicides to Surface and Ground Water — *R.C. Warner*
- Vapor Pressure Deficit Control as an Alternative Environment Control Technique for Plant Production — *R.S. Gates*
- Variable-rate Seeding: A Cooperative Study with Kentucky Corn Producers — *S.A. Shearer*

Entomology

Research in Entomology has two major goals: 1) understanding biology of insects and related arthropods and their interactions with plants and animals and 2) development and implementation of safe and effective management tactics and strategies for pest species. Three broad areas of research are pursued, ranging from fundamental studies to practical field trials. Highlights of accomplishments for 1996 are reported here.

Behavior, Biosystematics, Ecology, and Evolution

- A previously unknown form of *Pandora* sp., an entomopathogenic fungus, was discovered. This finding suggests an entirely new (soil-borne) aspect of the ecology of this pathogen.
- More than 40 compounds induced by Japanese beetle feeding were identified from preferred host plants. Complexity of a plant's volatile blend seems to be an important cue used by this pest for host location. Constituents of leaf surface waxes play a role in resistance of some woody plants.
- Abundant pollen during sweet corn pollination led to higher densities of the predator *Coleomegilla maculata*, which feeds on pollen as well as insect pests.
- Populations of the predator *Coleomegilla maculata* were higher in sweet corn than in field corn; the predators developed faster feeding on sweet corn pollen than on field corn pollen.
- There is heritable variation in the pheromone blend of the black cutworm moth in populations from Kentucky, Egypt, and France. This finding indicates that the pheromone communication system could evolve in response to selection.
- Pheromones and sex attractants have now been identified for all of the prey species of a bolas spider, *Mastophora hutchinsoni*. This spider mimics the pheromones produced by female moths, and thus it attracts and captures male prey.
- The parasitoid complex of the introduced pine sawfly, *Diprion similis*, was characterized.
- In *Harmonia axyridis*, an introduced ladybird beetle, the extent to which diet affects development, size, and color depends on the genetic makeup of individuals. Successful colonizing and continued population growth of *H. axyridis* in Kentucky may therefore be facilitated by genetic variation underlying traits related to feeding, growth, survival, and reproduction.
- In *Nauphoeta cinerea*, male-male social interactions depend on the quantity of different pheromone components

that have conflicting roles. Increasing one component can eliminate the effects of increasing the other two components.

- Keys, descriptions, host records, and distribution records for 41 parasitic wasps of the braconid subfamily *Agathidinae* of Japan were published, including fifteen species new to science. The publication will allow biocontrol investigators to tap the reserve of Japanese species that are of potential use against introduced pests. To date, five species of *Agathidinae* have been introduced to the United States in classical biological control programs.
- Spider predation lowers densities of one group of forest-floor detritivores, but does not affect several other insects. Information on prey palatability and spider foraging behavior aids in predicting the actual impact of spider predation.

Pest Management and Applied Ecology

- Early-season release of *Pandora* sp. (generated from its overwintering form) was as effective at controlling tobacco aphids as conventional chemical controls.
- An improved method of incorporating biological control agents into economic decision tools was developed that can be easily implemented into traditional pest management programs.
- Field studies of white grubs and black cutworms in turfgrass revealed ways that cultural practices, including mowing, clipping removal, pH modification, top dressing, and resistant grasses, can be used to manage these pests with reduced reliance on insecticides.
- Studies of the horned oak gall, a severe pest of pin oak trees on horse farms, revealed periods when the wasps can be controlled with horticultural oils or other nontoxic means, thus safeguarding horses in adjacent pastures.
- A system was developed by which thrips and aphids on greenhouse crops can be controlled by overnight exposure to modified atmospheric conditions, without use of pesticides.
- The spider *Clubiona abbotii* consumed 20 to 25 corn earworm eggs per day on intact soybean plants when egg densities were high (32 to 64 eggs per plant).
- The predator *Nabis roseipennis* survived better and produced heavier adults when fed corn earworm eggs than when fed corn earworm larvae throughout its development.
- A newly identified third pheromone component of the black cutworm moth, (Z)-11-hexadecenyl acetate, improves trap efficiency by several fold. This finding should allow for more effective detection and monitoring of this pest.

- Evaluation of an expert system for management of potato leafhopper on alfalfa showed that this pest must be controlled at early regrowth of alfalfa after harvest to prevent stunting.
- Synthetic pyrethroid insecticides at rates as low as 0.0125 lb ai/acre gave excellent control of alfalfa weevil larvae and potato leafhopper on alfalfa.
- Autodissemination of the SeMNPV baculovirus for suppression of beet armyworm populations was investigated.
- Field studies support the efficacy of a new termite bait being marketed to homeowners in Kentucky. Of 22 chronically-infested houses baited in 1995, 18 (82%) exhibited no further signs of termite activity the following year.
- 350 nurseries were inspected and licensed in 1996, and 664 nursery dealers were licensed to sell nursery stock in the state.
- A third generation of permethrin analog was found to be effective against cattle horn flies as well as face flies. Organophosphate insecticides formulated into ear tags are still effective against horn flies but are ineffective against face flies.
- Preliminary field experiments demonstrated that predation by spiders and ground beetles may lower densities of Colorado potato beetles on potatoes and spotted cucumber beetles on cucumbers and squash.

Physiology, Genetics, and Molecular Biology

- A teratocyte (extraembryonic cell of a parasitic insect) gene that inhibits protein synthesis at the level of translation was isolated and expressed.
- A new type of hydrolase that acts on oxygen-nitrogen bonds was characterized from the midgut of the tobacco budworm.
- Regulation of dietary glucose for trehalose and glycogen synthesis in the tobacco hornworm was shown to be altered by parasitism.
- Quantitative genetic studies reveal that genetic variation among individuals influences the growth, reproduction, color, and development of the introduced ladybird beetle, *Harmonia axyridis*.
- Differences in the chemical composition of the pheromone of *Nauphoeta cinerea* are due primarily to genetic differences among individuals.
- Polydnavirus genomes appear to be segmented to regulate viral gene copy number and the level of viral gene expression; this finding provides insights into the evolutionary processes that produced segmented viral genomes.
- A teratocyte gene that inhibits protein synthesis at the level of translation was isolated and expressed. This activity is linked to the inhibition of growth in parasitized insects.

Research Projects

- Biological Control of Selected Arthropod Pests and Weeds — *K.V. Yeargan*
- Biology and Management of Insects Attacking Urban Landscape Plants — *D.A. Potter*
- Control Processes in a Terrestrial Food Web: Trophic Interactions of a Generalist Predator — *D.H. Wise*
- Cooperative Agricultural Pest Survey: Gypsy Moth — *B.C. Pass, P.M. Dillon*
- Damage Thresholds, Risk Assessment and Environmentally Compatible Management Tactics for White Grub Pests of Turfgrass — *D.A. Potter, K.F. Haynes, A.J. Powell*
- Determination of Resistance of Woody Landscape Plants to the Japanese Beetle — *D.A. Potter, T.R. Kemp, P.A. Weston, R.E. McNeil*
- Development and Integration of Entomopathogens into Pest Management Systems — *G.C. Brown*
- Development of Sustainable IPM Strategies for Soybean Arthropod Pests — *K.V. Yeargan*
- Ecology and Management of European Corn Borer and Other Stalk Boring Lepidoptera — *G.C. Brown*
- Evaluation of Controlled Atmosphere Anoxia Treatments as a Potential Disinfestation Technique for Thrips and Spider Mites in Greenhouses — *D.A. Potter, R.G. Anderson*
- Evaluation of Southern Region Integrated Pest Management — *B.C. Pass*
- Evolutionary Genetics of Developmental and Age-related Changes in Social Signals — *A.J. Moore*
- Impacts of Spiders in Food Webs of Crop and Forest Floor Ecosystems — *D.H. Wise*
- Insect Stress: Multitrophic Interactions between Parasites, Pathogens, and Allelochemicals — *D.L. Dahlman*
- Integrated Management of Arthropod Pests of Livestock and Poultry — *F.W. Knapp*
- Isolation of Biologically Active Secretory Products from an Endoparasite — *D.L. Dahlman*
- Kentucky's Agricultural Pesticide Impact Assessment Program — *B.C. Pass, M.P. Johnson, L.M. Unger*
- Mate Choice and Offspring Fitness — *A.J. Moore*
- Molecular Dissection of Polydnavirus Functional Activities — *B.A. Webb*
- Nursery Inspections — *B.C. Pass, J.T. Collins, C.W. Harper*
- Pathogenicity, Transmission, and Introduction of a Cytoplasmic Polyhedrosis Virus to Fall Webworm — *L.K. Reiske-Kinney*
- Pesticide Applicator Training 1996 — *L.H. Townsend*
- Real and Apparent Complexity in Polydnavirus Genomes — *B.A. Webb*
- Redundancy in Chemical Communication: Evolution of Sex Pheromone Blends — *K.F. Haynes, A.J. Moore*
- Semiochemical Mediation of Reproductive Behaviors in Moths — *K.F. Haynes*
- Sexual Selection and Plasticity in Social Behavior and Signals — *A.J. Moore, K.F. Haynes*
- Simple Dynamical Models for Incorporating Biological Control Agents into IPM Decision Making — *G.C. Brown*
- Spatial Dynamics of Leafhopper Pests and Their Management on Alfalfa — *B.C. Pass, J.C. Parr*
- The Role of Fluctuating Asymmetry in Sexual Selection — *A.J. Moore*
- Translation Factors from a Parasitic Wasp and Its Virus — *B.W. Webb*

Forestry

National, regional, and local interest in the future of forest resources stresses the need for strong research programs to provide information from which sustainable forest policies can be developed for these resources. These needs are further emphasized by conflicting social, economic, and political directions. Kentucky's 12.7 million acres of forest land (50% of its total land area) support important water supplies, habitat for diverse species, as well as important economic industries in forest products and recreation/tourism. The research mission of the Department of Forestry is to support wise management policies by conducting creative and rigorous studies addressing questions on forest ecosystem function and on the products of these ecosystems.

Dynamic programming has been used to model timber stand dynamics and optimal management prescriptions and has been expanded to incorporate site conditions and price fluctuations. Using this approach, improvement thinnings of shortleaf pine and yellow-poplar have been shown to increase economic returns by 5% to 15%. However, depending on market conditions, management for fiber may be preferable to sawtimber. These results suggest that management opportunities for landowners may be greater than previously considered.

There is increasing interest in obtaining accurate resource inventories in order to set effective forest policy. A temporal inventory data model has been developed which provides better process modeling, i.e., reconstruction of long-term forest transactions (growth, mortality, and timber sales). However, there are significant costs associated with such an inventory model in terms of database size, access, and management costs. These costs must be recognized as an important component of developing effective long-range forest policy.

A regional survey of forest ecosystem quality has been initiated using soil invertebrates as measures of ecosystem integrity and quality. Soil invertebrates respond quickly to changes in soil environment and may reflect long-term historical impacts on a forest site. Maintenance of ecosystem quality is a key component of ensuring sustained forest productivity.

Sediment production from forest management (especially timber harvesting and road construction) may have a significant impact on environmental quality. A runoff generation experimental field site has been established at Robinson Forest to characterize the distribution and importance of runoff generation mechanisms in the forested landscape. This study will characterize the natural production of runoff and sedi-

ment yield as well as production from human-induced sources, such as timber harvesting. A second experimental watershed has been established to monitor the effectiveness of Best Management Practices in preventing accelerated erosion from forest management activities.

Long-term studies of threatened and endangered species of forest-dwelling bats have determined basic biological requirements for survival and have helped establish management guidelines to ensure conservation of these species. Data on population dynamics, food preference and foraging habits, and roost selection have helped determine especially sensitive forest areas for bat conservation. Land management agencies, such as the Daniel Boone National Forest, have used this information to revise forest management policy and timber harvesting practices.

Studies have been initiated to evaluate the effects of timber harvest on genetic diversity of white oak populations of eastern Kentucky. Timber harvest may influence gene frequencies directly, or may impact mating system characteristics of residual trees and pollen and seed migration. Understanding these relationships may assist design of timber harvesting systems which will ensure sustainability of genetic diversity of this important timber species.

Observed losses of cations from the forest floor of northern hardwood forests have suggested possibly serious consequences for long-term productivity of these forests. Studies have been conducted in a variety of forest conditions to elucidate the cause of such losses. Cation loss was not common across stands of differing ages, as would be expected if acid deposition were the cause; nor was cation loss greatest in most rapidly growing stands, as might be expected if rapid nutrient accumulation in biomass of young stands following harvest were the cause. These results suggest that regional pollution and forest successional patterns cannot explain the observed losses.

Geographic positioning systems using satellite triangulation to determine ground location are among the new technologies being applied to forest management. Computation of area of GPS-traversed lands customarily uses the traditional coordinate method. However, no attempt has been made to examine new area determinations that might have distinctive computational benefits. Recent investigations have revealed that a numerical method, called Areas of Oriented Figures, may have the computational advantage of treating surveying data of convex and non-convex figures of many sides. Efficient algorithms are being developed to enable computer implementation of this numerical method.

Research Projects

- Development of Conservation Strategies for Forest-dwelling Wildlife Dependent on Topographic Habitat Features — *M.J. Lacki*
Economic Assessment of Surface Mine Reclamation Alternatives — *J.M. Ringe, M.H. Pelkki*
Effects of Forest Management Practices on Forest Nutrient Status — *M.A. Arthur*
Effects of State Interval Size and Number of Predictors on Dynamic Programming Solutions in Forestry — *M.H. Pelkki*

- Evaluating Differential GPS Positioning Accuracies in Forestry Applications — *C.J. Liu*
Indicators of Ecosystem "Quality" in the Mixed Mesophytic Forest — *P.J. Kalisz*
Intraspecific Phylogeography of Plant Mitochondrial DNA — *D.B. Wagner*
Roost Selection of Bats in Forests in Eastern Kentucky — *M.J. Lacki*
Sources of Runoff and Sediment Production in Forested Watersheds — *D.L. Brown*

Horticulture and Landscape Architecture

The Horticulture and Landscape Architecture faculty conduct mission-oriented research, combining basic and applied approaches related to the production and use of horticultural crops and landscape architecture. Research activities are concentrated in the broad topics of integrated crop management, crop improvement, environmental regulation of plant growth and development, and landscape architecture.

Integrated Crop Management

The Controlled Water Table irrigation system consists of a capillary mat placed on a horizontal surface with one end of the mat suspended vertically in a small trough of water or nutrient solution. The amount of water, in a pot of growing medium sitting on the horizontal surface, is determined by the vertical distance from the water to the bottom of a pot. A liquid level controller maintains the water level in the trough. As plants remove water from the growing medium, it is replaced immediately by water from the trough. The system may be used in commercial greenhouse production and residential homes.

A novel approach to rooting cuttings under mist uses capillary mat subirrigation. The capillary mat reduces the amount of water in the rooting medium during misting cycles and prevents the medium from drying during the night when the mist is off. This provides a better environment for root development.

Nursery crop production system research revealed that composted yard trash is an effective container medium component when comprising 25% of a pine bark medium volume. Media comprised of 50% to 100% composted yard trash were not acceptable for outdoor production of two holly species. A three-year field nursery fertilization study revealed that banded application of 250 pounds of nitrogen per acre per year, split into spring and fall applications, is adequate for the production of the seven landscape plants tested. Field herbicide trials resulted in expanded herbicide labels for nursery crops and offer hope for the formulation of herbicides with a broad spectrum of control and tolerance for landscape use.

A study of wholesale prices for selected landscape plants over a 12-year period revealed that annual price increases were fairly constant from 1980 until 1989. A drastic drop in price increases that occurred in the 1989-90 catalog year was attributed to the severe drought of 1988. Price increases rebounded in the 1990-1991 year and leveled off the following years.

Three promising resistant bell pepper varieties were identified after the second year of testing 20 cultivars at two lo-

cations under induced and natural epidemic bacterial leaf spot conditions. Rapid adoption of resistant pepper cultivars occurred following recommendations from our 1995 trials; no losses occurred in growers' fields with resistant varieties compared to 50% losses in fields of susceptible varieties at the same localities.

Crop Improvement

The widespread occurrence of genes and transcripts for the ¹⁴C-N-methyltransferase responsible for the methylation of Lysyl residue 14 in the large subunit of Rubisco is present in many plant species. Methyltransferase has an exceptionally high affinity for Rubisco and represents the first demonstration of a protein with a tight and specific interaction with the world's most abundant and important plant protein responsible for the fixation of CO₂ by all green plants. The characterization of this protein and gene has resulted in the identification of a molecular vehicle capable of the selective delivery of positive agents to Rubisco. The significance of this protein and its activity, as well as its potential usefulness in the molecular engineering of higher yielding crop species, is being explored.

The activity of the key enzyme for sorbitol metabolism, sorbitol dehydrogenase, in apple fruit is induced by sorbitol, indicating that the source-sink interaction in apple is more dynamic than previously thought. Reduced sorbitol production resulting from defoliation due to disease and insect problems may also create "weaker" fruit, leading to poor quality at harvest.

The human dietary component E-2-hexenal, a natural constituent of all higher plants, was found to be the most effective volatile compound evaluated for controlling microbial pathogens on plant-derived foods such as strawberries and table grapes. Collaborative studies were undertaken to determine the mode of action of this compound and to establish effectiveness in inhibiting bacteria growth. Due to their natural origin and metabolism by the fruit, the volatile compounds may offer a biologically safe, consumer-friendly solution for eliminating postharvest losses due to pathogens such as gray mold.

Somatic embryos developed in legume species often show severe abnormalities and are difficult to convert to seedlings. A system was developed using eastern redbud to initiate adventitious shoots from malformed somatic embryos. Multiple shoots were initiated on each somatic embryo that could be subsequently rooted and converted to plantlets. This system bypasses the problem with germination of malformed somatic embryos and could be applicable to other legume species.

Seed germination in vinca (*Catharanthus roseus*) progresses in a typical triphasic pattern for fresh weight gain. Ethylene evolution was not observed in vinca seeds until radicle emergence (48 hrs) even though detectable levels of 1-aminocyclopropane-1-carboxylic acid (ACC) were present during this period. ACC oxidase activity was not detectable until 48 hours and preceded in vivo ethylene production. Our data suggest that ACC oxidase is a key regulatory step in the biosynthesis of ethylene during early stages of germination and that ethylene has a role in radicle growth.

How arthropods perceive plant-borne chemical repellents is an important aspect of plant-insect interaction. *Lycopersicon hirsutum* is a wild species of tomato that is highly defended from arthropods. Certain genotypes produce repellents in trichome secretions. For the three chemical repellents tested (2,3-dihydrofarnesoate, 2-pentadecanone and laurate), most quantitative behaviors were similar except average distance from the stimulus. For the two acids, distance from the stimulus tended to decrease the longer mites were exposed, indicating that the stimulus to leave becomes stronger as time passes. Data suggest that 2-pentadecanone may act as an attractant when perceived by olfaction, but as a repellent when perceived by touch.

Long-term cultivar and rootstock research is making significant contributions to the Kentucky fruit industry. Blueberry cultivars in descending order of yield (kg/bush) in the second year were Duke (2.2), Sierra (1.9), Bluecrop and Patriot (1.8), Toro (1.5), Blue Gold (1.4), Nelson (1.1), and Sunrise (0.7). The table grape cultivars, in descending order of yield (kg/vine) were Niagara (14.8), Concord (10.5), Moored (8.1), Mars (7.6), Golden Muscat (3.5), Captivator (3.0), Reliance (2.6), Challenger (2.4), Canadice and Himrod (0.25), and Glenora (0.0). The 1990 apple cultivar/rootstock planting is trained to the Dutch slender spindle system and supported by electrical conduit fastened to a wire trellis. Early production via this system allows growers to quickly establish orchards with newer, more profitable cultivars. The semi-dwarf apple rootstock trial utilizes the French vertical axe system. The mortality of trees in this study was higher on M.26 (40% survival) than on the other five rootstocks (98% survival).

Sorgoleone, a long-chain hydroquinone, is present in large quantities (greater than 80%) in most root exudates of various sorghum accessions. It is a potent inhibitor of seedling growth, oxygen evolution, and electron flow in photosystem II. It exhibits similar potent inhibitory activity within PSII as diuron or metribuzin and demonstrates similar competitive binding characteristics within the Q_b binding pocket. Also, sorgoleone's binding and 3D configuration in the Q_b niche are similar to that of other PSII inhibitors such as lentacil and metribuzin. This is attributed to sorgoleone's quinone moiety and in particular its unique electrostatic charge distribution. Apparently sorgoleone can persist in soil systems for days to weeks, causing inhibition of plant growth.

Environmental Regulation of Plant Growth and Development

Commercial quality cut roses were produced in a single stem production system from single node cuttings. The initial node of origin was significantly correlated to the final stem length of the single stem 'Lady Diana' rose and to the number of days for axillary bud break on the cutting. More than 70% of the cut stems harvested from these cuttings were 46 to 75 cm long. An adaptive neuro-fuzzy inference system was built to predict the rose status and to make critical environmental control decisions based on the principle of economic optimization.

Landscape Architecture

A Geographic Information System (GIS) database has been developed for land use planning in the 28 counties comprising the eastern Kentucky coalfields. A major effort in 1996 was made in analyzing institutional impediments to reuse of devastated lands, particularly in regard to legal/regulatory requirements and interpretations of these in the field. The database will be available through Kentucky Cooperative Extension in 1997.

As part of a restoration/preservation study of downtowns, models for organizing the relationship of parking and pedestrian movement to downtown destinations have been identified, and a method for evaluating pedestrians' circulation experiences has been developed. Recommendations and design criteria will be conveyed as conceptual unifying principles and guidelines and as examples of applications through case study descriptions.

Research Projects

- All-American Selection Trial Garden for Annual Flowering Plants — *S. Bale*
- Antimicrobial Properties of Naturally Occurring Volatile Compounds from Plants — *T.R. Kemp*
- Blackberry and Raspberry Cultivar Evaluation — *G.R. Brown and D. Wolfe*
- Composted Yard Trash as a Container Medium Component — *D.L. Ingram and B.R. Roach*
- Control of Root Outgrowth by Copper Hydroxide in Capillary Mat Plug Production — *R. Geneve and J. Buxton*
- Daylily Cultivar and Production System Evaluation — *W. Dunwell*
- Densities and Secretions of Trichomes on *Lycopersicon* — *J.C. Snyder*
- Developmental and Environmental Influences on Carbohydrate Partitioning in Fruit Crops — *D.D. Archbold*
- Downtown Morphology and Circulation Potential — *N. Crankshaw*
- Evaluation of Scab-immune Apple Cultivars Using IPM and Organic Techniques — *J.G. Strang*
- Evaluation of Cut Flower Species for Adaptability to Improved Greenhouse Production Practices and Extended Postharvest Life — *R.G. Anderson*
- Evaluation of Bacterial Leaf Spot-resistant Pepper Cultivars and Breeding Lines in Epidemic and Disease-free Environments — *B. Rowell, R.T. Jones, and W. Nesmith*

- Evaluation of Controlled Atmosphere Anoxia Treatments as a Potential Disinfection Technique for Thrips and Spider Mites in the Greenhouse — *R.G. Anderson, D.A. Potter, and R.S. Gates*
- Fall Vegetable Greenhouse Production — *D. Spalding, R. Anderson, and B. Rowell*
- Fertilization Systems for Field-grown Nursery Crops — *D.L. Ingram*
- Field Performance of Herbaceous Perennial Plants — *W. Dunwell*
- Herbicide Evaluations in Vegetable Crops and Woody and Herbaceous Ornamentals — *L.A. Weston and R.E. McNeil*
- Integrated Crop Management Program for Apples — *J. Hartman, G.R. Brown, D. Wolfe, and R. Bessin*
- Interaction between Ethylene and Polyamines during Seed Germination and Early Seedling Growth — *R.L. Geneve*
- Knowledge-based Manufacturing System for Optimization of Greenhouse Rose Production — *R.G. Anderson and R.S. Gates*
- Light, Temperature and CO₂ Effects on Carbohydrate Metabolism in Bedding Plant Seedlings — *J.W. Buxton*
- Mechanism and Significance of Post-translational Modifications in the Large Subunit of Ribulose Bisphosphate Carboxylase/Oxygenase — *R.L. Houtz*
- Monitoring Wild Ginseng in Kentucky — *R.T. Jones*
- Peach Tree Cultivar Evaluation for Hardiness — *J.G. Strang and D.D. Archbold*
- Planning Model for the Development of Mountaintop Removal-valley Fill Mining Sites — *T.J. Nieman*
- Post-translational Modifications in Ribulose Bisphosphate Carboxylase/Oxygenase — *R.L. Houtz*
- Pot-in-pot Tree Production Practices for Kentucky — *R.E. McNeil*
- Producing and Marketing Deciduous Holly Berries for Use in Floral Designs — *R.T. Jones, M. Witt, and S. Bale*
- Production Research on Artificial Shade and Woods-grown Ginseng and Goldenseal — *R.T. Jones*
- Rootstock and Interstem Effects on Pome and Stone Fruit Trees — *G.R. Brown*
- Seedless Table Grape Cultivar and Training System Evaluation — *G.R. Brown and D. Wolfe*
- Southeast Tree Fruit Cultivar Evaluation — *G.R. Brown, D. Wolfe, J. Strang, and R.T. Jones*
- Supersweet Corn Cultivar Evaluations — *J.G. Strang, R.T. Jones, J. Snyder, D. Slone and D. Lowry*
- Sustainable N Management: Intensive Crop Production and Improved Water Quality — *D.C. Ditsch, R.T. Jones, R.C. Pearce, and J.H. Grove*
- Technical and Economical Efficiencies of Producing and Marketing Landscape Plants — *R.E. McNeil*
- The Physiological Bases for Allelopathic Interference of Sorghum spp. as Mediated by Sorgoleone — *L.A. Weston, C.I. Nimal, and J.S. Pyrek*
- The Landscape of Main Street: An Assessment of the Urban Landscapes of Kentucky's Main Street Towns — *N. Crankshaw*
- Use of Natural Volatile Compounds for Control of Microbial Spoilage and Quality of Strawberry during Modified Atmosphere Storage — *D.D. Archbold, T.R. Kemp, B. Langlois, and M. Barth*
- Use of the Tomcast Model for Disease Management in Processing Tomatoes — *B. Rowell, R. Bessin, and W. Nesmith*
- Using Available Tobacco Resources to Diversify Farm Enterprises and Increase Income — *R.T. Jones and D.C. Ditsch*

Nutrition and Food Science

The Department of Nutrition and Food Science is supported by the Agricultural Experiment Station at the University of Kentucky through supplies and salary of four faculty members and one chairperson, each with joint appointments and approved projects in the AES. AES faculty perform extensive research, administrative, service, and instructional assignments for the Department of Nutrition and Food Science.

The main thrust of AES-supported departmental research includes nutrition in relation to eating disorders, cardiovascular disease, cancer, nutrient-drug interactions, and alcohol. All faculty were invited speakers at national and/or international research meetings. All faculty reviewed manuscripts and grant applications for journals and funding agencies. The faculty is committed to research, service, and instruction and will continue to carry out their duties in accordance with the guidelines and goals of the University of Kentucky, the College of Human Environmental Sciences, and the College of Agriculture.

Alcohol

Chen

Acetaminophen or acetaminophen-containing drugs are commonly used analgesics and antipyretics. Tylenol, the most popular preparation of acetaminophen in the U.S., can be obtained without prescription. Although acetaminophen is a safe drug at the therapeutic dose, overdose or long-term use can cause chronic hepatitis or necrosis. The mechanism by which acetaminophen produces hepatotoxicity remains unclear. It has been suggested that oxidative stress may be a possible mechanism. Thus, the possible protective effects of antioxidants and glutathione (GSH) precursors against the APAP-induced alterations of antioxidant defense enzymes in hepatocyte culture were studied. Rat hepatocyte culture plates were divided into five groups and pretreated with none, (α -tocopheryl succinate (TS), L-ascorbic acid (AA), N-acetylcysteine (NAC), or S-adenosyl methionine (SAM). One half of the plates in each group were treated with acetaminophen. The cells were harvested and homogenized, and biochemical assays were performed. Pretreatment with each of the four compounds partially protected against APAP-induced damage of the integrity of the cell membrane, partially prevented the decline of cellular GSH levels, and completely prevented the depression of glutathione reductase (GR) activity. The results suggest that vitamin E, vitamin C, or GSH precursors can lessen APAP toxicity through modulation of APAP-induced depression of antioxidant defenses.

Cancer

Chow

The toxic effect of endotoxin is partly attributable to increased generation of reactive oxygen species by endogenous mediators of inflammations. While a number of studies have shown that endotoxemia is associated with increased generation of lipid peroxidation products and altered activities of antioxidant enzymes, relatively little is known concerning the effect of endotoxin treatment on the status of small molecular weight antioxidants and if protein oxidation occurs during endotoxemia. Acute effect of endotoxin on the generation of oxidation products and status of small molecular weight antioxidants was, therefore, examined in mice. Male B6C3 mice were treated with 0, 20, 40 or 80 mg of endotoxin/kg BW, i.p., for 24 hours. Liver, heart, and lung were measured for the levels of malondialdehyde (MDA), conjugated dienes, protein carbonyls (PC), glutathione (GSH), ascorbic acid (AA), retinol, and tocopherols. Relative to the vehicle controls, endotoxin-treated mice had significantly ($p < 0.05$) higher levels of MDA and CD in the liver and heart of endotoxin-treated mice. The levels of PC were not significantly altered by endotoxin in liver, heart, or lung. Endotoxin-treatment resulted in lower levels of GSH in the liver, heart, and lung, of retinol in the liver and heart, and of alpha-tocopherol in the liver, while the levels of AA were higher in the liver and heart.

Cancer

Glauert

The overall project goals are to determine the mechanisms of peroxisome proliferator-induced liver carcinogenesis and to determine the role of dietary antioxidants. The following research accomplishments occurred in 1996:

1. We developed a transgenic mouse which has increased (3 to 4 fold) liver-specific expression of the antioxidant enzyme catalase. This will allow us to determine the role of active oxygen in the promotion of liver carcinogenesis by various agents;
2. We demonstrated that eicosanoids and the peroxisome proliferator ciprofibrate are co-mitogenic in cultured rat hepatocytes. This implies that specific eicosanoids must be present for peroxisome proliferators to exert possibly their promoting effects;

3. We found that transcription factor NF- κ B is activated by the peroxisome proliferator ciprofibrate and by phenobarbital in the liver, which may provide a new mechanism by which these agents exert their tumor-promoting activity. Since NF- κ B is an active oxygen-activated transcription factor, dietary intervention (e.g., vitamin E, selenium, synthetic antioxidants) may prevent the activation of this transcription factor.

Cardiovascular Disease

Hennig

Kentuckians are experiencing a high incidence of nutrition-related health problems, which may be due to overconsumption of fat and lack of protective nutrients such as antioxidants. Zinc, for example, is an essential component of biomembranes and is necessary for maintenance of membrane structure and function. There is evidence that diet-derived zinc can provide antiatherogenic properties by preventing metabolic physiologic derangements of the vascular endothelium. Because of its antioxidant and membrane-stabilizing properties, zinc appears to be crucial for the protection against cell-destabilizing agents such as polyunsaturated lipids and inflammatory cytokines. Zinc also may be antiatherogenic by interfering with signaling pathways involved in apoptosis. Most importantly, we have evidence that zinc can protect against inflammatory cytokine-mediated activation of oxidative stress-responsive transcription factor, such as NF- κ B and AP-1. It is very likely that certain lipids and zinc deficiency may potentiate the cytokine-mediated inflammatory response and endothelial cell dysfunction in atherosclerosis. Thus, the antiatherogenic role of zinc appears to be in its ability to inhibit oxidative stress-responsive factors involved in disruption of endothelial integrity and atherosclerosis.

Eating Disorder

Mercer

We have accomplished several goals producing new findings placing the central H1 receptor in the hierarchy of regulation of food intake: the central H1 receptor is involved in physiological responses to nutrient-imbalanced and restricted intake diets; food intake is inversely proportional to central H1 concentration in male rats fed diets varying in energy content, protein content, and protein quality; female rats normally have higher H1 receptors than male rats; diet composition affects H1 receptor levels in a gender-specific manner; bioperiodicity is present in central H1 receptor trafficking; parameters of periodicity are gender-specific and modified by diet.

Preliminary findings in our lab carried out on animal tissues received from other laboratories indicates that:

- tumor-bearing mice have increased H1 receptors (consistent with anorexia);
- cigarette-smoking rats have increased H1 receptors (consistent with weight loss);
- Zucker obese rats have impaired H1 response to dietary manipulation (consistent with obesity).

Hypothalamic tissues from humans obtained at autopsy indicate bioperiodicity of H1 receptors.

Research Projects

- Dietary Vitamin E/Fat and Oxidative Damage — C. Chow
- Lipid-mediated Endothelial Injury — B. Hennig
- Mechanism of Hepatocarcinogenesis by Peroxisome Proliferators and Inhibition of Dietary Antioxidants — H. Glauert
- Nutrient-alcohol/Oxidative Drug Interaction — L. Chen
- The Histaminergic System and Eating Disorders — L.P. Mercer

Plant Pathology

Healthy plants are key to sustainable agriculture and a secure food base for the Commonwealth. Plant pathology is an integrative scientific discipline which ranges from the whole plant, to interrelationships in the microbial world, to mechanisms of cellular dysfunction in the disease process. Although the department's research endeavors focus particularly on fungal and viral pathogens, attention is also paid to disease problems created by bacteria, nematodes, and non-living causal agents.

Mycology

Fungi are a constant threat to production agriculture and demand scientific study toward their management. Some pertinent departmental projects are summarized below.

Research on mycorrhizal stunt disease of tobacco is being extended to determine if pathogenic mycorrhizal fungi are involved in the general phenomenon of yield depression associated with continuous cropping without rotation. Systems are being developed to reduce the necessity of having to conduct initial field experiments to detect such yield declines. The approach is to devise greenhouse techniques to mimic field experiments. The approximately two dozen mycorrhizal fungi present in a field soil are isolated. After several cycles of continuous or rotated growth of crop plants, the mycorrhizal fungal communities are again determined and evaluated for crop response.

A long-term research program was initiated to investigate stalk rot of field corn, the primary disease problem of corn in Kentucky. Stalk rot is difficult to control by conventional means because of the extreme variability in incidence and severity between years and locations. The long-term goals of the research are to better understand the genetics and physiology of anthracnose stalk rot of corn and to develop sustainable methods for control of this and other stalk rot diseases.

In order to study the interactions between two related fungi superinfecting a grass plant, two hygromycin-resistant (hygR) transformants of the choke disease pathogen *Epichloe typhina* were introduced into perennial ryegrass seedlings which already possessed the related mutualistic, seed-transmissible symbiont, *Neotyphodium lolii* (= *Acremonium lolii*). Forty-six plants superinfected with *E. typhina* and *N. lolii* were vernalized and tested for seed production, transmission of the fungi in seeds, and hygR of the transmitted fungi. In no instance was *E. typhina* infection sustained through vernalization and seed set, but the original *N. lolii* infections remained stable. Based on these results, *N. lolii* may be able to protect host plants from choke disease.

The first pathway-specific step of ergot alkaloid biosynthesis in the ergot fungus *Claviceps purpurea* is catalyzed by 4-(dimethylallyl) tryptophan (DMAT) synthase. A gene, *dmaW*, thought to encode this enzyme has been cloned. The function of this gene was tested and results confirmed that *dmaW* encoded authentic DMAT synthase. The cloned *dmaW* was used as a probe to identify clones of putative *dmaW* from *Neotyphodium coenophialum* (= *Acremonium coenophialum*). Tall fescue infected with this endophyte has enhanced fitness in some environments. The putative *N. coenophialum dmaW* encoded a polypeptide with a 121 amino acid region of similarity to *C. purpurea* DMAT synthase.

Evidence was developed suggesting that polygalacturonase (PG) produced by the chestnut blight fungus *Cryphonectria parasitica* may be a virulence factor. When the gene encoding a basic PG produced in culture by *C. parasitica* was disrupted, however, the disrupted mutant caused cankers similar in size to those induced by the nondisrupted virulent parental strain. The basic PG produced in culture is therefore not required for virulence. Additional PGs, acidic rather than basic, are produced in cankers, and research is in progress to determine if these are required for virulence.

A multidisciplinary research effort culminated in demonstrating that failed stand establishment in alfalfa seeded in spring is often due to the root rot fungus *Aphanomyces euteiches*. As a result of this research effort, the University now recommends only *Aphanomyces*-resistant alfalfa varieties for spring seedings.

Apple scab conidial inoculum dose was positively correlated with infection rates, especially at the lower end of its temperature range for infection.

Research showed convincing proof that *Pyricularia grisea* is causing increasing, sometimes catastrophic, loss of high-maintenance perennial ryegrass golf fairways and provided strong direction for the development of control strategies.

Virology

Plant pathologists discovered viruses some 100 years ago, setting the stage for a new field of endeavor in the agricultural and medical sciences. Certain of the major research emphases in the department concerning viruses are highlighted below.

The virus genes responsible for differential host responses were studied in two systems: the response of Tabasco pepper to tobacco etch virus (TEV) and the response of the tobacco cultivar TN86 to a resistance-breaking strain of tobacco vein mottling virus (TVMV). By construction of chimeric viruses, two regions of the TEV genome and a single region of the TVMV genome were shown to be involved in the responses.

The importance of retention of virus particles in the stylets of aphids to the transmission of potyviruses by these insects was confirmed. The ability of aphids to transmit TEV was abolished or drastically reduced by probing oil-sprayed membranes or leaves prior to virus acquisition, and by probing oil-sprayed leaves after virus acquisition. The lack of transmission correlated with inability to retain virus particles in the stylets. Fasted aphids transmitted TEV more efficiently than unfasted aphids. Results indicated that fasting may eliminate a plant component(s) which interferes with the retention of virus in the stylet food canal.

The NIB protein of TMV was expressed in *Escherichia coli* as a GST-fusion protein or a his6-tagged protein. His6-NIB is able to undergo nonspecific binding to the 3'-untranslated region of TMV RNA in the absence of other proteins. The fusion protein possesses the poly (U) polymerase activity that is a hallmark of picornavirus-encoded polymerases. In addition, the protein is able to use full-length TMV RNA as a template for RNA synthesis. A fusion protein containing a mutation in the highly conserved GDD motif had only 7% of the activity of the wild type protein. Mutants of TMV in which the tyrosine residue that links the NIa-VPg to the viral RNA were constructed. There was no evidence of accumulation of progeny viral RNA in plants or protoplasts inoculated with mutant transcripts.

A search for the first region in TMV RNA to be encapsidated by the coat protein during the assembly of virus particles was undertaken. Protoplasts were collected at various times after they had been inoculated. TMV and the fragments of progeny viral RNA that were protected by coat protein subunits were isolated. The first fragments were detected in protoplasts that had been infected for 35 to 40 minutes, and accumulation of completely assembled progeny virions was noted after 45 minutes.

The dsRNA genome of *Helminthosporium victoriae* 190S tobitivirus (Hv109SV) was shown to consist of two large overlapping open reading frames (ORFs). The 5' proximal ORF codes for the capsid protein (CP) and the 3' ORF for an RNA-dependent RNA polymerase. Although the capsid of Hv190SV is encoded by a single gene, it is comprised of two major closely related polypeptides, p88 and p78. Whereas p88 is a phosphoprotein, p78 is nonphosphorylated. The finding that p78 and p88 share a common N-terminal amino acid sequence is consistent with the determination that N-terminal, but not C-terminal, CP deletions were incompetent for assembly. Evidence was obtained that p78 is derived from p88 via proteolytic cleavage.

Strains of peanut stunt cucumovirus (PSV) were classified into two distinct subgroups, I and II, based on Western and Northern blot analyses using antisera and cloned cDNA probes to strains PSV-ER and PSV-W. These results were corroborated by nucleotide sequence analyses of full length cDNA clones of RNA3 from representative strains of the two subgroups.

Disease Management

The department pursues a variety of studies to implement research findings for plant disease control, several of which are briefly described below.

Tobacco production was hit hard by a blue mold epidemic in 1996, with production losses in excess of \$150 million. However, without appropriate disease management practices, the consequences of blue mold to the tobacco industry might have been catastrophic. The North American and Kentucky Blue Mold Warning Systems were coordinated, and the emergency labeling of Acrobat MZ was organized through the department. Long-term monitoring of the changing sensitivity of the blue mold fungus to Ridomil was instrumental in convincing the Environmental Protection Agency (EPA) that Ridomil was no longer a reliable control, allowing alternative fungicide use for this crisis situation. Likewise, evaluation of experimental fungicides for blue mold control under greenhouse, plant bed and field conditions provided data vital to EPA's granting of emergency-use fungicide treatment. Field and greenhouse evaluations were made of the sensitivities of current burley cultivars to blue mold.

Black shank is a perennial problem to tobacco growers. About 15 pre-plant soil fumigation test plots were monitored over the past two growing seasons. The results obtained from this research indicated the need to recommend pre-plant soil fumigation at problem sites, particularly in light of the recent sharp decline in crop rotation. Other current production practices contributing to increased black shank were also determined.

Efficacy of fungicides for management of dogwood powdery mildew, which has become a more common disease in recent seasons, was assessed.

Efforts to determine the effects of various cropping sequences on soybean cyst nematode (SCN) population densities and associated soybean yields from a multi-year study indicated that a race shift had likely been initiated in the presence of resistant soybean varieties. In most cases, growing corn following the resistant soybean cultivars tested reduced SCN levels.

Research Projects

- Altering Ergot Alkaloid Biosynthesis by the *Acremonium* Endophyte of Tall Fescue — *C.L. Schardl*
- Basis of Virus Virulence/Host Resistance in Potyvirus-tobacco Interactions — *T.P. Pirone*
- Biological Improvement of Chestnut and Management of the Chestnut Blight Fungus — *L. Shain*
- Characterization of Viral Genes and Gene Products Which Mediate Aphid Transmission and Cell-to-cell Movement — *T.P. Pirone*
- Chemical Controls for Tobacco Diseases under Greenhouse, Float Bed, Plant Bed, and Field Conditions — *W.C. Nesmith*
- Cultural Practices for Managing Spring Dead Spot of Bermudagrass — *P. Vincelli*
- Corn Stalk Rot Diseases — *L.J. Vaillancourt*
- Defining and Mapping the Genes of Caulimoviruses — *R.J. Shepherd*
- Disassembly and Early Gene Expression of RNA Plant Viruses — *J.G. Shaw*

Distribution of Bacterial Leaf Scorch in Kentucky — *J.R. Hartman*
 Effect of Conidial Inoculum Dose and Temperature on Development of Apple Scab — *J.R. Hartman*
 Effect of Cropping Sequence on Soybean Cyst Nematode Reproductive Ability and Soybean Yield — *D.E. Hershman*
 Effect of Reduced Tillage on Disease Development in Wheat — *D.E. Hershman*
 Evaluation of Disease Management Strategies for Tobacco and Vegetables — *W.C. Nesmith*
 Evaluation of Fungicides and Biocontrol Products for Control of Turfgrass Diseases — *P. Vincelli*
 Evaluation of Landscape Austrian Pines for Pine Tip Blight Disease and Assessment of Management Strategies — *J.R. Hartman*
 Forage Legume Viruses: Identification and Genetic Resistance for Improved Productivity — *S.A. Ghabrial*
 Host-pathogen Interactions between *Castanea* sp. and the Chestnut Blight Fungus — *L. Shain*
 Identification of Disease-resistant Cultivars of Turfgrasses and Corn — *P. Vincelli*
 Integrated Management Program for Sclerotinia Crown and Stem Rot of Alfalfa — *P. Vincelli*
 Interactions of Tobacco with the Biotrophic Fungus *Peronospora tabacina*: Potential for Enhanced Disease Resistance — *C.L. Schardl*
 Management of Rhizosphere Dynamics to Control Soilborne Pathogens and Promote Plant Productivity — *J.W. Hendrix*
 Molecular Basis of Disease in a Virus-infected Plant Pathogenic Fungus — *S.A. Ghabrial*
 Mycorrhizal Fungi in Relation to Health and Vigor of English Boxwood — *J.W. Hendrix*
 National Dogwood Anthracnose Survey — *J.R. Hartman*
 Natural and Modified Grass Endophytes as Agents for Biological Protection — *C.L. Schardl*
 Pest Predictive Technology for Apple Disease Management — *J.R. Hartman*
 Plant-fungal Endophyte Interactions: Potential for Cultivar Improvement in Species of *Festuca* and *Lolium* — *M.R. Siegel*
 Phylogenetics of *Epichloë* Species and Related Grass Mycosymbionts — *C.L. Schardl*
 Potyvirus Replication and Pathogenicity — *J.G. Shaw*
 Proteinaceous Inhibitors of *Cryphonectria parasitica* in Chestnut Bark — *L. Shain*
 Role of Coat Protein and Helper Component in Aphid Transmission of Potyviruses — *T.P. Pirone*
 Rotation of Resistance Genes to Maintain *Heterodera glycines* below the Damage Threshold — *D.E. Hershman*
 Structure and Function of the Viral dsRNAs of the Plant Pathogenic Fungus *Helminthosporium victoriae* — *S.A. Ghabrial*
 The Role of Specific Viral Genes and Gene Products in Potyviral Pathogenicity, Host Range, and Aphid Transmission — *J.G. Shaw and T.P. Pirone*
 Transformation of Plant Pathogenic and Plant Mutualistic Fungi — *C.L. Schardl*
 Transgenic Resistance to Bean Pod Mottle Virus and Soybean Mosaic Virus in Soybeans — *S.A. Ghabrial*
 Urban Tree and Ornamental Plantings Disease Evaluations — *J.R. Hartman*
 Viral Satellite RNAs as Riboregulators of Gene Expression in Tobacco — *S.A. Ghabrial*
 Wheat Seed Treatment and Foliar Fungicide Screening — *D.E. Hershman*

Regulatory Services

The Division of Regulatory Services administers state laws pertaining to the manufacturing, processing, labeling, and marketing of commercial feed, fertilizer, seed, and raw milk. Its purpose is to protect farmers and other consumers from poor quality, mislabeled, or misrepresented products and to protect agricultural businesses from unfair competition from those who might take short cuts in the quality of their products.

Feed, fertilizer, and seed are monitored in the manufacturing or retail channels for reasonable and acceptable compliance with state laws through label review, product inspection, and sampling and analyzing of products. Raw milk is monitored during marketing to assure an accurate and equitable exchange between producers and processors and to ensure the integrity of milk from farm to processor.

Ten regulatory inspectors and one auditor travel throughout the state, collecting samples, inspecting facilities, and auditing records. Additionally, full-time, temporary inspectors check and sample specialty feed, fertilizer, and seed products throughout the state.

The Division also offers to growers and homeowners seed testing, soil testing, poultry litter and animal manure testing, and water and nutrient solution analyses for greenhouse production and float-bed systems of seedling production.

Auditing Program

H.S. Spencer

Audits of sales and fee payments were made on 267 of 688 milk, seed, feed, and fertilizer businesses in Kentucky to verify check-off and tonnage fees. Fees assessed to help pay the costs of inspecting, sampling, and analyzing commodities in accordance with Kentucky laws are: fertilizer, 50 cents per ton; feed, 35 cents per ton; and seed, 4 to 24 cents per unit. During May, raw milk is assessed a check-off fee of 3 cents per 100 pounds. Although the tobacco seedling law was repealed in 1996, some fees were collected for 1995 in 1996.

Income from fees in 1996:

Feed	\$638,656
Fertilizer	616,340
Milk	55,045
Soil testing	149,353
Seed tags, testing, and licenses	357,141
Tobacco seedlings	22,151
TOTAL	\$1,838,686

These cash receivables were substantiated on 2,688 fertilizer tonnage reports, 3,092 feed tonnage reports, 792 seed reports, and 18 milk reports. These reports were checked for accuracy and compared with field audits of the firms submitting them. An additional \$3,000 was realized through auditing and correcting inaccurate reports.

Milk Regulatory Program

B.W. Cox

The milk regulatory program administers the Kentucky Creamery License Law, which establishes regulations for ensuring that dairy farmers are accurately paid for the milk they produce and that the integrity of milk is maintained from the farm to the processor. In 1996 the division:

- Reviewed and issued licenses to 13 milk buyers, 56 testers, 475 sampler-weighers, and 8 raw milk transfer stations.
- Analyzed and administered action on 9,000 official samples and checked 3,500 producer pay records.
- Conducted 64 inspections at 16 milk laboratories.
- Analyzed 1,000 exchange samples from commercial laboratories.
- Trained and examined 71 sampler-weighers and 11 testers.
- Conducted 500 inspections of 225 sampler-weighers.

Feed Regulatory Program

C.E. Miller

The feed regulatory program provides consumer protection for the purchasers of livestock feed and pet foods, maintains a marketplace environment that promotes fair and equitable competition for the feed industry, and helps ensure the safety and wholesomeness of animal products as human food. In 1996 the program:

- Administered actions on 4,434 official samples of commercial feed involving 26,126 official tests to monitor about 3 million tons of commercial mixed feed and feed ingredients distributed in Kentucky.
- Administered a cooperative program with the FDA on six feed mills that mix restricted drugs in feed. An additional 69 state inspections were conducted on mills that mix non-MFA drugs in feed to ensure compliance with medicated feed regulations. Thirty-one mills that mix no drugs were inspected to ensure compliance with labeling, manufacturing, and storage practices.

- Conducted 7,500 label reviews and maintained product registration for about 15,000 products from 900 companies.
- Cooperated with the Association of American Feed Control Officials in conducting the 32nd Annual Feed Management Seminar at Shakertown, Kentucky. Thirty-three professionals from 20 states, Canada, and the FDA attended.

Fertilizer Regulatory Program

D.L. Terry

The Kentucky Fertilizer Law ensures that fertilizers sold in Kentucky are clearly and accurately labeled so that consumers can make informed purchases of fertilizer and be assured of its quality. The law also protects the legitimate fertilizer industry from unfair competition. In 1996 the division:

- Administered actions on 4,606 official and 274 unofficial samples of fertilizer involving 13,277 tests of the 972,000 tons of fertilizer distributed in Kentucky.
- Reviewed labels and registered 3,995 products from 472 firms, including 240 who manufactured custom blends of fertilizers.

Inspection Program

F. Herald

The inspection program aims to achieve industry compliance with the consumer protection laws that the Division is charged with administering. This responsibility is carried out by inspectors strategically located throughout the state, each with an assigned area to cover. They inspect manufacturing plants, processing facilities, storage warehouses, and retail stores; collect official samples of feed, pet food, fertilizer, milk, and seed; review records; and offer advice and assistance to clientele in improving their operations to achieve compliance with the laws. In 1996:

- A team of nine inspectors performed 5,253 inspections of the processing, manufacturing, and marketing of feed, fertilizer, and seed. In addition, three temporary inspectors traveled throughout the state during March, April, and May to inspect and sample specialty feed, fertilizer, and seed products in urban areas, and one full-time permanent milk inspector covered the state to administer the Creamery License Law.

- Inspectors collected the following official samples for laboratory verification of appropriate constituents and quality:

Feed	4,434
Fertilizer	4,606
Seed	2,962
Milk	9,000

Seed Regulatory Program

D.T. Buckingham

The seed regulatory program assures Kentucky farmers of quality seed and promotes fair and equitable competition among seed dealers and seedsmen. In 1996 the program:

- Collected and tested 2,962 official seed samples.
- Issued stop-sale orders on 376 official seed samples and 192 violative seed lots at seed dealer and seed processor locations.
- Conducted a cooperative program with USDA, AMS on trueness-to-variety testing and on interstate shipments of seed.
- Issued 243 permits to label seed and registered 396 seed dealers.

Seed Testing Services

E.E. Fabrizius

The seed testing program provides the seed industry and seed growers of Kentucky with competent, reliable, and timely analyses of their seeds for labeling requirements and quality assurance. In 1996 the seed laboratory tested the following number of samples:

Official seed samples	2,962
Regular seed samples (service)	4,805
Certified seed samples (service)	428
Tobacco seed samples (service)	907
TOTAL	9,102

In addition to routine testing for purity, germination, and noxious weed seed, the laboratory offers testing for seed vigor (accelerated aging and cold test), seed moisture, seed size (seed count), and germination following fungicide application. The laboratory also provides tests for the presence of the fungal endophyte present in tall fescue seed and live plant tillers to seedsmen and livestock owners in the state.

Soil Testing Service

V.W. Case (Lexington)
D.L. Kirkland (UKREC, Princeton)

Soil testing provides farmers, homeowners, greenhouse operators, surface mine specialists, and others with scientific information about the fertility status of their soils. In partnership with the Cooperative Extension Service, it also provides them with lime and fertilizer recommendations based on soil tests. We also offer analyses of poultry litter and animal wastes for farmers and farm advisors, water and nutrient solution analyses for greenhouse operators and float-bed seedling producers, and non-routine, optional soil tests for University of Kentucky researchers.

Soil samples analyzed in 1996 were:

Agriculture	27,618
Home lawn and garden	6,200
Strip-mine reclamation	49
Commercial horticulture	627
Greenhouse	84
Research	8,567
TOTAL	45,480

The Lexington lab also analyzed 130 soil samples for triazine residue, 72 animal waste samples, and 187 water and nutrient solution samples for greenhouse and float-beds.

Robinson Substation

Research and demonstration trials at Quicksand are primarily conducted by the departments of Agronomy, Horticulture, and Plant Pathology. Agronomy research activities include livestock forage management, no-till corn production, tobacco nitrogen management, variety trials for corn, and the revegetation of surface-mined lands. Horticulture research consists of trials with fruit, vegetable, and ornamental cultivars for both commercial and home gardens and horticultural crop production on reclaimed surface-mined lands. Plant Pathology trials deal primarily with evaluating cultivar resistance in tobacco, peppers, and field corn.

Agronomy

Yield of corn from the long-term no-till plots continued to be good and were exceptionally high in 1996 due to the well-distributed 23 inches of rain received during the period from May to September. Nitrogen yield response in 1996 topped out at 218 bushels per acre with 160 lbs fertilizer nitrogen per acre. Average yield at 160 lbs nitrogen per acre over the 16 consecutive years of no-till corn production in this study is 169 bushels per acre. The range of grain yield for 160 lbs N/A over this time span has been 145 to 218 bushels per acre with yields of more than 170 bushels per acre in eight of the 16 years. Yields in 1996 showed little influence from three years of wheat cover crop as compared to not planting a wheat cover crop.

Soil test levels of phosphorus in the top three inches of soil have been built to and maintained at about 100 lbs per acre, but decrease to 20 to 30 lbs per acre in the 3- to 6-inch soil depth, and to below 20 in the 6- to 12- inch depth. This indicates that sustained high yields of corn can be obtained from surface applications of fertilizer phosphate on no-till fields without having to build up soil test phosphorus levels below the surface three inches of soil.

A study was initiated to evaluate the effectiveness of dormant herbicide applications compared to traditional foliar herbicides for control of kudzu. Kudzu is considered a noxious weed in many locations throughout eastern Kentucky. Dormant herbicide applications were applied in early April before new stem growth occurred for the season. Early evaluations in late May indicated that kudzu stem emergence was delayed when dicamba was applied as a dormant treatment. However, by late summer kudzu growth was equal to the untreated plots based on later evaluations. Best late-season control was obtained with dicamba and glyphosate applied as a foliar treatment in late May.

A 27-acre educational site is currently being developed on the Laurel Fork of the Robinson Forest to demonstrate the agricultural potential of reclaimed surface-mined land. Field trials addressing mined land revegetation for livestock forage production, wildlife habitat, and other herbaceous species/variety identification are available for evaluation and study. Production practices necessary for apple, blueberry, blackberry, and hydrangea production are also demonstrated at this site. In cooperation with Kentucky State University, utilization of irrigation ponds for rainbow trout and freshwater shrimp production is being monitored.

The Robinson Experiment Station serves as one of three primary testing sites for tall fescue, orchardgrass, and red clover varieties in Kentucky. This project is managed by the University of Kentucky Forage Variety Testing Program to assist forage producers in eastern Kentucky with the selection of forage species and varieties evaluated under more site-specific growing conditions, such as surface-mined lands.

The evaluation of pre-sidedress nitrogen tests on burley tobacco continued in 1996. The Robinson Experiment Station is one of several sites where this work has been conducted. Initial results indicated that a soil nitrate test just before sidedressing time had some value of predicting cured leaf yields. Further testing is being conducted to determine the feasibility of using the soil nitrate test to predict the need for additional sidedress nitrogen. The development of such a test could improve the efficiency of nitrogen use on burley tobacco.

In 1996 a demonstration float bed was constructed at the Robinson Experiment Station to promote new tobacco production technologies in eastern Kentucky. The float bed was used for both produce and agent training programs. Interest in float bed use has increased significantly in this region over the past two years.

Poultry litter is a source of plant nutrients, especially nitrogen (N), phosphorus (P), and potassium (K). With the growth of the poultry industry in Kentucky, more litter/waste has become available to grain producers. A field experiment was conducted to compare the availability of plant nutrients contained in litter to that of fertilizer sources in a no-till corn/soybean rotation.

The preliminary data suggest that poultry litter will provide considerable quantities of plant available P and K to the crop in the first year after application. Litter K appears to be fully available the first year, while only about 75% of the litter P is available in that season. To the extent that litter P and K were not removed by corn in the first season, they were available to the following soybean crop.

Horticulture

A summer annual flower cultivar evaluation was conducted as part of a statewide effort to determine which cultivars would develop well and produce attractive bloom and foliage for Kentucky's commercial and home landscapes. Forty annual flower cultivars were evaluated at Quicksand as part of this cooperative trial with Lexington and Princeton. This is the fourth year of annual testing at Quicksand, and several cultivars did well during a very wet year. Recommended annuals in 1996 include Dahlia "Calico Mix," Salvia, Impatiens "Blitz Salmon," Impatiens "Tempo White," Impatiens "Blitz Orange," Begonia "Varsity Rose," Begonia "Rio Scarlet," Celosia "Flamingo Feather," and Zinnia Linearis Orange.

The departments of Horticulture and Plant Pathology continued the field evaluation of 21 bell pepper cultivars with different levels of resistance to bacterial leaf spot (BLS). Nineteen cultivars with some BLS resistance and two susceptible check cultivars were screened against three races (BLS races 1, 2, 3) of the bacterial pathogen. A susceptible inoculated bell pepper cultivar (Merlin) was used as the inoculum source. Last year was an abnormally wet year, and conditions were very favorable for disease spread. Fruit of each pepper cultivar was evaluated not only for yield but for overall appearance. There were significant differences among the resistant bell pepper cultivars in yield, appearance, and BLS disease ratings. Cultivars with the highest marketable yields and quality were XPH 12205, Boynton Bell, and XR3 Aladdin. Other acceptable cultivars included PR 9300-8, Enterprise, and Commandant. Additional trials are planned for 1997.

Market demand for fall decorations resulted in an ornamental corn cultivar trial. Kentucky growers need attractive, highly marketable, large-eared (8-12 in.) cultivars. Nine large-eared ornamental corn cultivars were evaluated in 1996. One cultivar, Earth Shades Indian, produced large, primarily dark red ears that were very attractive and did not exhibit the lodging problems seen in other cultivars. Previously, Rainbow and Fiesta were recommended for commercial use. Cultivar trials in 1997 will evaluate three new hybrid large-eared ornamental corn cultivars in hopes of finding cultivars with bright colors, tight shucks, excellent tip fill, and resistance to lodging. Prior to 1997, all large-eared ornamental corn cultivars were open pollinated. Results from these trials have been used to help expand sales of Kentucky's fall ornamental crop acreage by providing a more diverse sales package.

A study involving fall cole crops as scavengers of residual fertilizer nitrogen was jointly conducted by Agronomy and Horticulture faculty to evaluate the use of fall cole crops (i.e., cabbage and/or broccoli) to recover residual fertilizer N following high-value summer annual cash crops. Summer crops, tobacco and sweet corn, were managed according to University of Kentucky recommendations and received a fertilizer N application of 300 lb N/A and 148 lb N/A, respec-

tively. In 1996, immediately following the harvest of the summer annuals, broccoli, winter rye, and fallow treatments were established. Fertilizer N was applied to the broccoli at rates of 0, 50, 100, and 150 lb N/A. A broccoli yield response to 50 lb N/A was measured in 1996. A cabbage yield response to 50 lb N/A was measured in 1994, whereas no cabbage yield response to fertilizer N was measured in 1993 or 1995. Residual N following sweet corn and tobacco was slightly lower in 1996 and 1994 than in 1993 and 1995, indicating that a yield response to fertilizer N could be expected when residual soil $\text{NO}_3\text{-N}$ levels fall below 20 ppm. Soil nitrate-N and plant total N uptake were also measured to determine the usefulness of broccoli (compared to winter rye and fallow) to recover residual N and minimize the potential for N loss to the environment. Fall broccoli or cabbage has been found to be as effective as rye in capturing nitrogen that would normally be lost due to leaching. Additional research in 1997 is planned using broccoli again as the fall nitrogen scavenger.

Work on field diagnostic tools for estimating nitrogen needs for fresh market tomatoes is another ongoing cooperative project between Agronomy and Horticulture. Hand-held Cardy meters were found to be effective for rapidly measuring fresh petiole sap nitrate nitrogen levels in fertigated fresh market tomatoes. The Cardy meter was also effective in determining soil nitrate - N levels prior to planting. Four nitrogen (N) fertigation rates were tested, including a 0 N/A rate. Results in 1996 showed that there was a significant US#1 tomato yield increase due to fertilizer nitrogen application, even when an initial soil mineral N test showed 70 lb N/A available for plant growth. 1996 was a very cool, wet year, and plant root growth may have been limited more than in previous years. Early season Cardy meter petiole sap $\text{NO}_3\text{-N}$ levels temporarily dropped below levels considered optimum for tomato plant growth and yield. The 0 lb/A nitrogen treatment was the slowest to return to optimum levels, whereas the 225 lb/A N treatment was the fastest. Later in the season, petiole sap nitrate nitrogen readings were at adequate levels for all treatments. Previous trials have shown that many Kentucky soils contain high levels of plant available nitrogen prior to fertilization. However, under very cool, wet conditions, early season supplemental (extra) nitrogen may be needed to maintain optimum yields when soil tests indicate that adequate levels of mineral N are not available. Relating soil mineral nitrogen level to tomato N response will eventually enable growers to customize their fertilizer nitrogen applications to specific growing sites and weather conditions.

Work will continue in an effort to determine the correlation between weather, initial soil mineral nitrogen levels, and tomato crop nitrogen requirements. Preliminary petiole sap NO_3 data appears to be less variable and may be a better predictor of tomato yield response than soil tests. Additional tomato nitrogen fertility trials are planned for 1997.

Research Projects

- Annual Flower Cultivar Evaluations — *S. Bale and R.T. Jones*
Bacterial Leaf Spot-resistant Bell Pepper Cultivar Evaluations —
B. Rowell, R.T. Jones, and W.C. Nesmith
Determining Fertilizer Nitrogen Needs for Fresh Market Tomatoes —
R.T. Jones and D.C. Ditsch
Establishing the Value of the Phosphorus and Potassium Contained in
Poultry Litter for a No-till Corn and Soybean Rotation — *J.H.
Grove*
Evaluation of Soil and Crop Diagnostic Tools to Improve Nitrogen
Management for Burley Tobacco — *R.C. Pearce and D.C. Ditsch*
Fall Cole Crops as Scavengers of Residual Fertilizer Nitrogen — *D.C.
Ditsch, R.T. Jones, and R.C. Pearce*

- Floral Hydrangea Cultivar Evaluation — *R.T. Jones, S. Bale, and
R. McNeil*
High Bush Blueberry Cultivar Evaluation — *R.T. Jones, J. Strang, and
J. Brown*
Kudzu Control Study — *J.D. Green and D.C. Ditsch*
Ornamental Corn Cultivar Evaluations — *R.T. Jones and Mary Witt*
Response of No-till Corn to Carryover Soil N Recovered by a Winter
Wheat Cover Crop — *K.L. Wells, D.C. Ditsch, and W.O. Thom*
Tall Fescue Adaptability as Affected by Endophyte Fungus Infection
— *T.D. Phillips, D.C. Ditsch, L.M. Lauriault, M. Collins, and J.C.
Henning*
Tall Fescue, Orchardgrass and Red Clover Variety Trials — *L.M.
Lauriault, J.C. Henning, and D.C. Ditsch*

Rural Sociology

The Rural Sociology research program is designed to advance the development and understanding of social structure and social processes in the changing rural society. Through an integrated program of research, extension programs, and instruction, the faculty focus on initiatives which are organized around two major areas: sociology of agriculture and community and human resource development. Within each major area, several substantive areas of specialized activity exist.

Community and Human Resource Development

Social and Economic Factors Affecting Family Well-Being. Rural society's prospects for growth and development ultimately rest with the capacities of people. Human resources refer to the skills, abilities, and understanding required for people to function effectively in families, the workplace, organizations, and communities. Specific areas of study are rural health, youth, family dynamics and functioning, leadership development, aging, and the role of service agencies in human resource development.

Findings from a study of women's home-based work indicate that home-based businesses are often a hidden segment of a local economy. The impacts of home-based businesses on the local economy depend on the characteristics of the rural women operating these businesses, the types of economic activities themselves, and the personal and family consequences of operating the businesses.

Faculty continue working with the Kentucky Kids Count Consortium to produce an annual data book highlighting various indicators of the demographic and socioeconomic status of Kentucky's children and families. Ninety-seven focus groups of families and service care providers have determined social and economic factors influencing family well-being which will be used to guide Kentucky's state plan for family preservation and family support services. The evaluation of extension programs with youth at risk, exploring school and family factors that influence educational achievement and the ability of communities to retain youth, has been conducted.

Other research results on factors affecting family well-being have had important impacts in Kentucky. Faculty have presented results through commissions and publications on the status of families and the state and local social service structures that affect families. These results have been used for legislative recommendations by the Kentucky State Legislative Research Commission on Poverty and for developing social service plans by the Kentucky Cabinet for Human Resources, Department of Social Services.

Community Resource and Rural Development. Communities are the settings in which people interact with the larger society to provide basic human services, employment, and solutions for local problems. Within the sphere of rural development policy, the local community is the context for initiation of economic and social development. Research findings have led to policy proposals based on the concept that programs can gain local legitimacy by sharing policy authority with local citizens. Local rural development efforts would be led by local boards which involve a broad base of local citizens but with federal funding to facilitate implementation of local plans.

Sociology of Agriculture

Trends in Agriculture. As an example of research regarding changing agriculture, findings show that Kentucky's dairy industry is confronting both a changing marketplace and changing conditions of production. Research results have been used to determine educational needs of these farmers for planning and adapting their operations to meet future conditions.

Kentucky is part of a regional study designed to assess how an increasingly globalized agro-food system impacts commodity producers in the region and the state. Changes in commodity programs will be examined as a result of increasing globalization of agricultural markets, as well as how domestic fiscal concerns may impact farm enterprise structure and rural community well-being in rural regions of the South and of Kentucky.

Other research programs are examining the impacts of changes in the international trade system on the structure of agriculture and rural communities, impacts upon Kentucky rural communities from changes in the federal tobacco program and potential decreases in income from tobacco farming, and post-productionist agricultural sector adjustment strategies that emphasize the production of "social goods," such as environmental amenities and landscape preservation, in lieu of traditional commodity production objectives.

Intergenerational Changes in Farm Family Perspectives of the Environment. Environmental and natural resource issues are critical components of rural development policy. Program focus is on natural resource utilization and economic development, impacts of natural resource production, and relationships among environmental factors and quality of life. Findings have led to the development of a conceptual framework of how cultural groups transform the nonhuman environment symbolically and thereby mesh the social, cultural, and nonhuman environments into structures of symbols and beliefs that are shared among members of the group.

Research Projects

Customer Service Initiative of Kentucky Cabinet of Natural Resources and Environmental Protection — *T. Greider*

Family Preservation and Family Support Services Evaluation and Technical Assistance — *P. Dyk and L. Garkovich*

Forest Fire in Eastern Kentucky: How the Role of Fire in the Ecosystem Is Perceived and Controlled — *L. Garkovich*

Harlan Youth Employability Program Evaluation — *P. Dyk*

Home-based Work: The Relationship of Informal Employment to Health among Mid-life and Older Rural Women — *L. Garkovich*

Impacts of Structural Change in the Dairy Industry — *L. Garkovich*

Intergenerational Changes in Farm Family Perspectives on the Environment — *T. Greider*

Kentucky Kids Count — *L. Garkovich and G. Hansen*

Ohio River Valley Development — *L. Swanson*

Reexamining East Asian Land Reform: Class and Culture in Action — *L. Burmeister*

Rural Communities Initiative of the Rural Policy Research Institute — *L. Swanson*

Rural Restructuring: Causes and Consequences of Globalized Agricultural and Natural Resource Systems — *L. Burmeister*

Service Sector Penetration and Household Earning Processes: A Comparative Study of Rural Communities — *R. Harris*

Social and Economic Factors Affecting the Well-being of Kentucky Rural Families — *P. Dyk, R. Harris, L. Garkovich, and G. Hansen*

Social Capital Attributes of Families, Schools, and Communities — *P. Dyk*

Veterinary Science

Maintenance of the health and promotion of the welfare of Kentucky's animal industry continue to be primary concerns of the Department of Veterinary Science. The department's three functional units, the Livestock Disease Diagnostic Center, Maxwell H. Gluck Equine Research Center, and the Equine Blood Typing and Research Laboratory, help support the needs of the state's livestock, horse and companion animal industries through a comprehensive range of nationally and internationally recognized diagnostic, extension, and research programs. Publication of the *Equine Disease Quarterly* continues with the January 1997 issue distributed to 11,400 individuals in 70 countries. In the second year since its inception, six foreign students participated in the Kentucky International Thoroughbred Institute (KITI) program, an equine training program for international students which provides hands-on experience and education in farm and racetrack management.

Maxwell H. Gluck Equine Research Center

The Gluck Center continues to consolidate its reputation, both nationally and internationally, for the quality and success of its programs in selected fields of equine research. Infectious diseases, immunogenetics, immunoparasitology, pharmacology, and reproductive physiology remain the areas of primary research emphasis of the program.

Immunogenetics—A gene map of the horse is being constructed in collaboration with various other laboratories both in the U.S. and throughout the world. The major emphasis is on identifying genetic markers to place on the map and mapping genes to specific chromosomes. In the past year, the gene for combined immunodeficiency disease in Arabian horses was located on horse chromosome 9, thereby confirming identification of a candidate gene for the disease. Other ongoing studies are focused on characterization of the major histocompatibility gene family which controls the immune response to certain infectious and noninfectious diseases.

Infectious Diseases—Infectious diseases remains an area of critical significance for both breeding and racing industries not only nationally, but also with respect to the international movement of horses, semen, or equine embryos.

As in the last few years, research on equine herpesvirus 1 has focused on better defining the immune correlates of protection in the horse to this economically important viral infection. In a collaborative study with the Animal Health Trust, Newmarket, in the United Kingdom, virus-specific cytotoxic T-lymphocytes are being investigated for their role in protecting against abortion caused by the virus. This work will

improve our understanding of how to achieve more effective protection against this ubiquitous virus in the horse.

Several significant accomplishments have emerged from the research program on equine influenza. Certainly, the most notable has been the finding that strains of equine-2 virus in the eastern and western hemispheres have evolved independently of one another with some evidence of cross-over. This has led directly to the international adoption of new recommendations for appropriate virus strains representative of each lineage to be included in updated equine influenza vaccines. In collaboration with the equine pharmacology program, the pharmacokinetics of the anti-influenza virus drug, rimantadine, have been established for the horse, and a dosing protocol has been determined which can ameliorate the clinical signs of influenza in horses.

The equine lentivirus that causes equine infectious anemia is an important cause of morbidity and mortality in equids worldwide. Research on equine infectious anemia virus has important implications for equids, and as a model for HIV infection in humans. Our attempts to define the determinants of virulence in this disease have been aided by the generation of an infectious pathogenic clone of the virus through site-directed mutagenesis. Recent research on developing a vaccine to protect equids against infection with equine infectious anemia virus would suggest that multiple viral determinants may be necessary for stimulating sterile immunity against this disease.

Ongoing collaborative studies with the University of California-Davis have helped to better define the location and nature of the neutralization determinants on the genome of equine arteritis virus. In conjunction with researchers at the Animal Diseases Research Institute, Lethbridge, Canada, the RT-PCR assay has been validated for use as a diagnostic test for detection of the carrier state in the stallion. Studies of natural outbreaks of EVA have furthered our understanding of the pathogenicity of semen isolates of EAV in terms of abortion in mares and illness and loss in neonatal foals.

The equine rotavirus vaccine, which received a conditional USDA license permitting its use in the state of Kentucky at the beginning of 1996, is now licensed for use in other states.

Mapping of the protective M protein (SeM) of *Streptococcus equi* has revealed numerous epitopes reactive with equine serum and mucosal antibodies on the N terminal half of the molecule. Great variability in specific epitope responses has been observed among individual horses. During convalescence from *S. equi* infection, the dominant serum antibody response to SeM is IgG. Strong mucosal IgA and IgG responses also occur. Commercial vaccines against strangles have been shown to elicit specific serum IgG responses but

not a mucosal IgA response. A PCR-based technique has been developed and validated for the rapid detection of *S. equi* in nasal swabs from infected horses.

Six proteins of *Leptospira kennewicki* immunoreactive for the horse have been partially characterized. One of these proteins, Lk90, isolated from the surface of *L. kennewicki* has been sequenced and expressed in *E. coli*.

Parasitology—A major health problem in yearlings with associated mortality was shown to be related to larval stages of small strongyles encysted in the large intestinal lining. Transmission patterns for natural infections with endoparasites were determined in lambs born on a farm in Kentucky in 1993 and 1994; *Haemonchus contortus* was the predominant species incriminated in both years.

Equine protozoal myeloencephalitis (EPM) was reproduced in normal horses by oral dosage with *Sarcocystis* sp. sporocysts from feral opossums. The experimentally infected horses developed *S. neurona*-specific antibodies in serum and cerebrospinal fluid which were detected by immunoblot analysis. Microscopic lesions consistent with EPM were observed in the CNS of experimentally infected horses. Aliquots of the sporocyst stock administered to horses produced sarcocystosis when fed to budgerigars and sparrows. This suggests that *S. neurona*, the etiologic agent of EPM, may be *S. falcatula* or a closely related *Sarcocystis* sp.

Pharmacology and Experimental Therapeutics—The Testing Integrity Program was brought on-line nationally in 1996, and a Testing Proficiency Program based on this initiative is under current development. The development of analytical tests for specific drugs or their metabolites has been assisted greatly by the acquisition of a Micromass Quattro-LC-Mg Mass Spectrophotometer. The Critical Metabolite Synthesis program has synthesized about fifteen important equine metabolites. The sample collection phase of the Musculoskeletal Injury Project in Kentucky was completed during this past year. In the area of new therapeutic approaches to diseases in the equine, we completed a project on endotoxic shock and are in the final stages of completing a research study on the potential value of amantadine and rimantadine for the treatment of equine influenza. A new therapeutic approach to equine protozoal myeloencephalitis was initiated in late 1996 which, hopefully, will provide a more effective, safer, and cheaper means of treating this economically important disease.

Reproductive Physiology—Considerable progress has been made in furthering our understanding of how the oviduct, uterus, and embryo communicate with each other biochemically to ensure a successful pregnancy in the mare. Work on differential gene expression and molecular cloning of gene products from embryonic tissues continues to advance our knowledge of how pregnancy is established and maintained in horses.

Research on the neuroendocrinological control of seasonal breeding in mares has revealed that the environmental cues which normally lead to cessation of the breeding season, though recognized by all mares, may be translated incorrectly in a small proportion of animals, resulting in continuation of

breeding activity throughout the year. The possibility that a metabolic cue by way of energy availability may play a pivotal role in this phenomenon is currently under investigation.

Livestock Disease Diagnostic Center

The Livestock Disease Diagnostic Center is a full-service laboratory providing a comprehensive range of diagnostic services in support of the livestock producers and veterinarians of Kentucky. An extension program for large ruminants, both beef and dairy cattle, was reinstated in early 1996. The following laboratory sections operate both independently and in support of the pathology program: virology, toxicology, serology/immunology, bacteriology, and clinical pathology. The number of pathology accessions increased by approximately 6% over that for the previous year. There were 67,498 diagnostic case submissions that included a total of 5,278 animals necropsied, making it one of the most heavily used veterinary medical diagnostic laboratories in the country.

During 1996, the Virology Section performed 10,121 tests to detect virus infections in a wide range of species together with 62,922 serologic tests to detect antibodies to various viruses. The different testing programs support diagnostic efforts to detect various infectious diseases, as well as regulatory and vaccine monitoring programs for the livestock and poultry producers of Kentucky.

Equine Blood Typing and Research Laboratory

A record number of blood samples (34,727) were tested for a record number of breed registries (30) by the Equine Blood Typing and Research Laboratory in 1996. Collaboration in the development of a gene map of the horse has been and will continue to be an area of major research focus of this program. Studies are in progress to detect genetic markers for disease traits of economic significance in certain species, e.g., *Epitheliogenesis imperfecta* in certain horse breeds, and juvenile cataracts in the Boston Terrier breed. Population genetic studies of the maintenance and significance of genetic variation in domestic and feral horse populations are continuing. A collaborative effort to develop a genetic map for the alpaca was initiated with researchers in Chile in 1996.

Research Projects

- A Study of the Prevalence and Distribution of Leptospirosis in the Horse Population in Central Kentucky and of the Significance of *Leptospira* as a Cause of Abortion in Mares and a Study to Determine Whether Certain Wildlife Species Can Serve as Long-term Reservoirs of the Organism on the Affected Farms — *K.B. Poonacha*
- Amantadine/rimantadine Prophylaxis of Equine Influenza — *T. Tobin*
- Analytical Methods Development — *T. Tobin*
- Application of New Technology for the Diagnosis of Equine Infectious Anemia — *C. Issel*
- Cell Culture Models for Analysis of Equine Influenza Viral (Host Discrimination) — *T.M. Chambers*
- Chemotherapeutic Control and Prevalence of Natural Infections of

- Internal Parasites of Equids — *E.T. Lyons*
Control and Transmission of Internal Parasites of Ruminants — *E.T. Lyons*
Critical Metabolite Synthesis — *T. Tobin*
Determining the Importance of Coronavirus in Respiratory Infections of Cattle — *M.L. Vickers*
Development of a Rapid Screening Test for EVA Antibodies — *M.L. Vickers*
Diagnostics for Equine Infectious Anemia — *C. Issel*
Differential Gene Expression During Early Equine Conceptus Development — *K.J. McDowell*
EIAV Vaccine Development — *C. Issel*
Epitope (B Cell, T Cell) Map of the Protective M Protein (SeM) of *Streptococcus Equi* — *J.F. Timoney*
Equine Protozoal Myeloencephalitis: Development of an Antemortem Diagnostic Probe and Epizootiologic Analysis — *D.E. Granstrom*
Equine Protozoal Myeloencephalitis: Isolation and Characterization of the Etiological Agent — *D.E. Granstrom*
Equine Protozoal Myeloencephalitis: Prevention and Control — *D.E. Granstrom*
Equine Respiratory Tract Mucosal Antibody Production in Response to Influenza Infection or Vaccination — *T.M. Chambers*
Functional Characterization of Immunogenic Proteins of *Leptospira Kennewicki* — *J.F. Timoney*
Gene Map of the Horse — *E. Bailey*
Genetic and Mechanistic Analysis of the Equine MX Antiviral Gene — *T.M. Chambers*
Genetic Basis of *Epitheliogenesis Imperfecta* in Saddlebred Horses — *E.G. Cothran*
Genetic Basis of Juvenile Cataracts in the Boston Terrier — *K.A. Graves*
Genetic Management of Feral Horse Populations on Public Lands — *E.G. Cothran*
Genetic Variation and Genetic Relationships of Populations of Domestic Horses — *E.G. Cothran*
Gonadotropin-releasing Hormone in Mares: Development of Methods for Measurement and Factors Affecting Secretion During Anestrus — *B.P. Fitzgerald*
Hemagglutinin Antigen of Equine-2 Influenza Virus: Surveillance and Molecular Characterization of Isolated Virus Strains — *T.M. Chambers*
How Conceptus Proteins Influence of Prostaglandin F_{2α} in Mare Endometrium — *K.J. McDowell*
Identification of the Equine Herpesvirus-1 Proteins and Their Respective Subregions that Elicit Cytotoxic T-lymphocyte Immune Responses in the Horse — *G.P. Allen*
Immunologic Management of Lentivirus Infections: EIAV — *C. Issel*
Immunological Significance of Hypervariable Region of M-like Protein of *Streptococcus Zooepidemicus* — *J.F. Timoney*
In Vivo Evaluation of Infectivity, Pathogenicity, and Immunogenicity of Different Novel Candidate Live Equine Influenza Virus Vaccines — *T.M. Chambers*
Mapping the Gene for Combined Immunodeficiency Disease in Horses — *E. Bailey*
Mechanism of Influenza-induced Cytopathology in the Respiratory Tract of the Horse — *T.M. Chambers*
Melatonin and Seasonal Reproduction in Mares — *B.P. Fitzgerald*
M-like Proteins of *Streptococcus Equi* in Virulence and Immunity — *J.F. Timoney*
Molecular Cloning and Characterization of Estrogen and Progesterone Receptors from Horse Endometrium: Comparisons between Pregnant and Nonpregnant Mares — *K.J. McDowell*
Molecular Cloning and Expression of Oxytocin in Equine Endometrium — *K.J. McDowell*
Mucosal Immunogenicity for the Horse of a Chimaera of Cholera Toxin and the Protective M Protein (SeM) of *Streptococcus Equi* — *J.F. Timoney*
Musculoskeletal Injury Program — *T. Tobin*
New Therapeutic Approaches to Equine Protozoal Myeloencephalitis (EPM) — *T. Tobin*
Pathogenesis of Natural Infection of Horses with *Sarcocystis Neurona* — *D.E. Granstrom*
Regulation of EIAV Replication and Disease — *C. Issel*
Research in Equine Pharmacology — *T. Tobin*
Sequence Analysis of Class I and Class II MHC Genes in Equidae — *E. Bailey*
Synteny Mapping of the Horse — *K.A. Graves*
Testing Integrity Program (TIP) — *T. Tobin*
The Development of Available Techniques for Immunohistochemical Detection of Viruses in Fixed Tissue — *M.L. Vickers*
The Effect of Electrolytes and Minerals on the Gastrointestinal Flora — *T.W. Swerczek*
Thresholds Research — *T. Tobin*
Use of Microencapsulation for Stimulation of Nasopharyngeal Mucosal Responses in the Horse — *J.F. Timoney*
Use of PCR Technology in the Diagnostic Laboratory — *M.L. Vickers*

Publications

All publication dates in this section are 1996 unless otherwise noted.

Annual Report

One Hundred and Eighth Annual Report of the Kentucky Agricultural Experiment Station. College of Agriculture, University of Kentucky, for 1995. C. Oran Little, Director. June.

Books and Book Chapters

Agricultural Economics

Francois-Poncet, J., H. Clemenson, D. Freshwater, and T. Rowley. *Partnerships in the Rural United States: A Review*. Paris, France: OECD.

Freshwater, D. Policy alternatives for stimulating rural employment. IN: R.D. Bollman, ed. *Rural Employment: An International Perspective*. London: CAB-International.

Marchant, Mary, ed. Trade and the Environment Conference Proceedings, American Agricultural Economics Association Annual Meeting, San Antonio, TX. 195 pages.

Agronomy

Ball, D.M., C.S. Hoveland, and G.D. Lacefield. *Southern Forages*. 2d ed. 264 pages. Norcross, GA: Potash and Phosphate Institute (PPI) and Foundation for Agronomic Research (FAR).

Bell, J.C., C.A. Butler, and J.A. Thompson. Soil hydrology and morphology of three Mollisol hydrosequences in Minnesota. IN: J. Wakely and S. Sprecher, eds. *Preliminary Investigations of Hydric Soil Hydrology and Morphology in Alaska, Indiana, North Dakota, Minnesota, and Oregon*. Wetlands Research Program Technical Report. U.S. Army Corps of Engineers, Waterways Experiment Station.

Collins, G.B., and R. Shepherd, eds. *Engineering Plants for Commercial Products and Applications*. 181 pages. Annals of the New York Academy of Sciences, Vol. 792. New York: New York Academy of Sciences.

Collins, M., and C.C. Sheaffer. Harvesting and storage of cool-season grass hay and silage. pp. 297-319. IN: L.E. Moser, D.R. Buxton, and M.D. Casler, eds. *Cool-Season Grasses*. Madison, WI: American Society of Agronomy.

Cornelius, P.L., J. Crossa, and M.S. Seyedsadr. Statistical tests and estimators of multiplicative models for cultivar trials. pp. 199-234. IN: M.S. Kang and H.G. Gauch, Jr., eds. *Genotype-by-Environment Interaction*. Boca Raton, FL: CRC Press.

Crossa, J., P.L. Cornelius, and M.S. Seyedsadr. Using the shifted multiplicative model cluster methods for crossover genotype-by-environment interaction. pp. 175-198. IN: M.S. Kang and H.G. Gauch, Jr., eds. *Genotype-by-Environment Interaction*. Boca Raton, FL: CRC Press.

Egli, D.B., and S.J. Crafts-Brandner. Soybean. pp. 595-623. IN: E. Zamski and A.A. Schaffer, eds. *Photoassimilate Distribution in Plants and Crops*. New York: Marcel Dekker, Inc.

Gardner, H.W., H. Takamura, D.F. Hildebrand, K.P.C. Croft, T.D. Simpson, and Y.P. Salch. Oxylin pathway in soybeans and its physiological significance. pp. 162-175. IN: G. Piazza, ed. *Lipoxygenase and Lipoxygenase Pathway Enzymes*. Champaign, IL: AOCS Press.

Hergert, G.W., W.L. Pan, D.R. Huggins, J.H. Grove, and T. R. Peck. The adequacy of current fertilizer recommendations for site-specific management. IN: F.J. Pierce, ed. *The State of Site-specific Soil Management*.

Hildebrand, D.F. Genetics of Soybean Lipoxygenases. pp. 33-38. IN: G. Piazza, ed. *Lipoxygenase and Lipoxygenase Pathway Enzymes*. Champaign, IL: AOCS Press.

Karathanasis, A.D., and B.F. Hajek. Elemental analysis by x-ray fluorescence spectroscopy. pp. 161-224. IN: D.L. Sparks, ed. *Methods of Soil Analysis*. 3rd ed. Madison, WI: Soil Science Society of America Monograph No. 9.

Maiti, I.B., and A.G. Hunt. Genetically engineered protection of plants against potyviruses. pp. 51-65. IN: P.M. Gresshoff, ed. *Current Topics in Plant Molecular Biology: Technology Transfer of Plant Biotechnology*. Boca Raton, FL: CRC Press.

Taylor, N.L., and K.H. Quesenberry. *Red Clover Science*. 226 pp. Dordrecht, The Netherlands: Kluwer Academic Publishers.

Thomas, G.W. Soil pH and soil acidity. pp. 475-490. IN: Methods of Soil Analysis. Part 3. *Chemical Methods*. Soil Science Society of America Book Series No. 5.

Animal Sciences

Bramley, A.J., L.S. Cullor, R.J. Erskine, L.K. Fox, R.J. Harmon, J.S. Hogan, S.C. Nickerson, S.P. Oliver, K.L. Smith, and L.M. Sordillo. *Current Concepts of Bovine Mastitis*. Madison, WI: National Mastitis Council.

Dawson, K.A., M.A. Rasmussen, and M.J. Allison. Digestive disorders and nutritional toxicity. Chapter 14. IN: P.N. Hobson and C.S. Stewart, eds. *The Rumen Microbial Ecosystem*, 2d ed. Hampshire, U.K.: Chapman and Hall Ltd.

Biosystems and Agricultural Engineering

Warner, R. Chapter 1: Sediment Control Basin Design and Construction; Chapter 2: Diversion Design and Construction; Chapter 4: Stream Channel Construction; Chapter 5: Vegetative Filters. IN: *Handbook of Western Reclamation Techniques*. Denver, CO: Office of Surface Mining.

Entomology

Potter, M.F. Termites. pp. 233-333. IN: S. Hedges, ed., *Handbook of Pest Control*. 8th ed. Cleveland, OH: Franzak & Foster Co.

Horticulture and Landscape Architecture

Geneve, R.L., J. Preece, and S. Merkle, eds. *Biotechnology of Ornamental Plants*. London: CAB International.

Hartmann, H.T., D.E. Kester, F.T. Davies, Jr., and R.L. Geneve. *Plant Propagation: Principles and Practices*. 6th ed. Englewood Cliffs, NJ: Prentice-Hall, Inc.

Nutrition and Food Science

Chow, C.K., V.L. Tatum, C.C. Yeh, M. Maynard, W. Ibrahim, G. Bruckner, G.A. Boissonneault, and C.B. Hong. Antioxidant function of carnosine, a natural histidine-containing dipeptide. pp. 155-161. IN: W.J. Xin and L. Packer, eds. *Natural Antioxidants: Molecular Mechanisms and Health Effects*. Champaign, IL: AOCS Press.

Glauert, H.P., J.T. Hong, L.K. Leung, and M.W. Wilson. The role of eicosanoid metabolism in carcinogenesis by peroxisome proliferators. IN: J.K. Reddy, T. Suga, G.P. Mannaerts, P.B. Lazarow, and S. Subramani, eds. *Peroxisomes: Biology and Role in Toxicology and Disease*. Annals of the New York Academy of Sciences, 804:719-721.

Nilakantan, V., Y. Li, B.T. Spear, and H.P. Glauert. Increased liver-specific expression of catalase in transgenic mice. IN: J.K. Reddy, T. Suga, G.P. Mannaerts, P.B. Lazarow, and S. Subramani, eds. *Peroxisomes: Biology and Role in Toxicology and Disease*. Annals of the New York Academy of Sciences, 804:542-553.

Plant Pathology

- Pirone, T.P., and S. Blanc. Helper-dependent transmission of plant viruses. 34:227-247. IN: *Annual Review of Phytopathology*.
- Schardl, C.L. *Epichloe* species: Fungal symbionts of grasses. 34:109-130. IN: *Annual Review of Phytopathology*.
- Schardl, C.L. Interactions of grasses with endophytic *Epichloë* species and hybrids. pp. 107-140. IN: G. Stacey and N. Keen, eds. *Plant-Microbe Interactions*. New York: Chapman & Hall, Publishers.
- Schardl, C.L., H.-F. Tsai, K.-R. Chung, A. Leuchtmann, and M.R. Siegel. Evolution of *Epichloë* species symbioses with grasses. pp. 541-546. IN: *Biology of Plant-Microbe Interactions*. St. Paul: International Society for Molecular Plant-Microbe Interactions.
- Siegel, M.R., and L.P. Bush. The role of defensive chemicals in grass fungal-endophyte associations. IN: J.T. Romero, P. T. Saunders, and P. Barbosa, eds. *Phytochemical Diversity and Redundancy in Ecological Interactions*. Recent Advances in Phytochemistry, 30:81-120.

Rural Sociology

- Luloff, A.E., and L.E. Swanson. Community agency and disaffection: Enhancing collective resources. Chapter 16. IN: L. Beaulieu and D. Mulkey, eds. *Investing in People: The Human Capital Needs of Rural America*. Boulder, CO: Westview Press.
- Swanson, L.E., R.P. Harris, J. Skees, and L. Williamson. 1995. African Americans in Southern rural regions: The importance of legacy. pp. 103-118. IN: J.B. Stewart and J.E. Allen-Smith, eds. *Blacks in Rural America*. New Brunswick: Transaction Publishers.
- Swanson, L.E. Social infrastructure and economic development. Chapter 7. IN: T. Rowley, D. Sears, G. Nelson, N. Reid, and M. Yetly, eds. *Rural Development Research: A Foundation for Policy*. Westport, CT: Greenwood Press.

Patents

Agronomy

- Evangelou, V.P. Pyrite silica coating. (U.S. and Canadian Patent Nos. 5,494,703 and 1,431,854, respectively.)

Biosystems and Agricultural Engineering

- Timmons, M.B., and R.S. Gates. Method for Controlling Environmental Conditions of Living Organisms Based upon Time-integrated Variables. U.S. Patent No. 5,573,179.

Progress Reports

- PR-384: 1995 Kentucky Vegetable Crop Research.
- PR-385: 1996 Agronomy Research Report.
- PR-386: 1996 Small Grains Variety Trials. C.R. Tutt, C.S. Swanson, D.A. Van Sanford and W.L. Pearce.
- PR-387: Kentucky Turfgrass Research, 1994-1995. A.J. Powell.
- PR-388: 1996 Kentucky Hybrid Corn Performance Test. W.L. Pearce, C.G. Poneleit, and J.E. Roberts.
- PR-389: 1996 Red Clover Report. L.M. Lauriault, J.C. Henning, N.L. Taylor, G.D. Lacefield, D.C. Ditsch, and R.E. Mundell.
- PR-390: 1996 Alfalfa Report. J.C. Henning, L.M. Lauriault, L.G. Brown, G.D. Lacefield, W.J. Peterson, P.C. Vincelli, and J.C. Parr.
- PR-391: 1996 Tall Fescue Report. L.M. Lauriault, J.C. Henning, T.D. Phillips, G.D. Lacefield, D.C. Ditsch, and E.L. Baker.
- PR-392: 1996 Orchardgrass Report. J.C. Henning, L.M. Lauriault, T.D. Phillips, G.D. Lacefield, D.C. Ditsch, and E.L. Baker.
- PR-393: 1996 Timothy Report. J.C. Henning, L.M. Lauriault, T.D. Phillips, D.C. Ditsch, and E.L. Baker.
- PR-394: 1996 Kentucky Soybean Performance Tests. E. C. Lacefield, C. Tutt, J.M. Wood, and T. Pfeiffer.
- PR-395: 1996 Beef Research Report

Regulatory Bulletins

- RB-268: *Commercial Feeds in Kentucky, 1995*. C.E. Miller.
- RB-269: *Seed Inspection Report, 1991-1995*. D.T. Buckingham.
- RB-270: *Analyses of Official Fertilizer Samples, July 1995-June 1996*. D.L. Terry.
- RB-271: *Seed Inspection Report, 1992-1996*. D.T. Buckingham.

Special Reports

- SR-96-1: *Nursery and Landscape Program Research Report*.
- SR-96-2: *Vegetable and Fruit Pest Management: A Survey of Pesticide Use and Other Pest Management Practices, 1994*. M.P. Johnson, J.G. Strang, S.C. Leak, S.M. Bailey.

Manuscripts Published in Refereed Journals

Agricultural Economics

- Cole, H., P. Kidd, S. Isaacs, M. Parshall, and T. Scharf. Difficult decisions: A simulation that illustrates cost effectiveness of farm safety behaviors. *Journal of Agromedicine*.
- Goetz, S.J., and D. Debertin. Rural population decline in the 1980s: Impacts of farm structure and federal farm programs. *American Journal of Agricultural Economics*, 78:3.
- Goetz, S.J., and D. Debertin. Rural-urban locational choices of medical doctors: A county-level analysis. *Review of Agricultural Economics*, 18(4):547-63.
- Goetz, S.J., D. Hu, and D. Debertin. A structural model of human capital and manufacturing sector change. *International Advances in Economic Research*, 2.
- Goetz, S.J., and D. Hu. Economic growth and human capital accumulation: Simultaneity and expanded convergence tests. *Economics Letters*, 51:355-62.
- Goetz, S.J., and D. Kemlage. TSD facilities location and environmental justice. *Review of Regional Studies*, 26.
- Goetz, S.J., and R. Ready. Economic growth vs. environmental conditions: A state-level analysis. *Growth and Change*, 27:97-110.
- Harrison, R.W., B. Bobst, F. Benson, and L. Meyer. Analysis of risk management properties of grazing contracts versus futures and options contracts. *Journal of Agricultural and Applied Economics*, 28(2):247-262.
- Infanger, C. Environmental regulatory reform and the unholy trinity: Unfunded mandates, risk assessment and property rights. *Journal of Applied Economics*, 28(1).
- Mawampanga, M., and D. Debertin. Choosing between alternative farming systems: An application of the analytic hierarchy process. *Review of Agricultural Economics*, 18(3):385-401.
- Munirathinam, R., M. Reed, and M. Marchant. International trade and foreign direct investment. *International Advances in Economic Research*.
- Ning, Y., and M. Marchant. Specification of a timber demand function: A case study for Douglas Fir Roundwood. *Kentucky Journal of Economics and Business*, 15:113-25.
- Ricks, D., and T. Woods. Strategic planning with horticultural crop industries. *Acta Horticulture*, 429:575-581.
- Ricks, D., and T. Woods. Strategic planning for improved competitiveness by regional commodity industries. *Journal of Food Distribution Research*.
- (In addition, members of the department published 46 abstracts.)
- #### Agronomy
- Back, K., and J. Chappell. Identifying functional domains within terpene cyclases using a domain-swapping strategy. *Proceedings, National Academy Science*, 93:6841-6845.
- Barbayanis, N., V.P. Evangelou, and V.Z. Keramidis. Potassium-ammonium-calcium quantity/intensity studies in the binary and ternary modes in two soils of micaceous mineralogy of Northern Greece. *Soil Science*, 161:716-724.

- Budziszewski, G., K.P.C. Croft, and D.F. Hildebrand. Uses of biotechnology in modifying plant lipids. *Lipids*, 31:557-569.
- Coyne, M.S. A cartoon history of soil microbiology. *Journal of Natural Resources and Life Sciences Education*, 25:30-36.
- Di, Rong, V. Purcell, G.B. Collins, and S. Ghabrial. Production of transgenic soybean lines expressing the bean pod mottle virus coat protein precursor gene. *Plant Cell Reports*, 15:746-750.
- do Valle Ribeiro, M.A.M., K.A. Gurney, and L.P. Bush. Endophyte *Acremonium lolii* in ecotypes of perennial ryegrass (*Lolium perenne* L.) collected in old Irish pastures. *Irish Journal of Agriculture and Food Research*, 35:151-157.
- Egli, D.B., and D.M. TeKrony. Seedbed conditions and prediction of field emergence of soybean seed. *Journal of Production Agriculture*, 9:365-370.
- Evangelou, V.P. Pyrite oxidation inhibition in coal waste by PO₄ and H₂O₂ pH-buffered pretreatment. *International Journal of Surface Mining, Reclamation and Environment*, 10:135-142.
- Fai, A.H.-T., and P.L. Cornelius. Approximate F-tests for multiple degree of freedom hypotheses in generalized last squares analyses of unbalanced split-plot experiments. *Journal of Statistical Computing and Simulation*, 54:363-378.
- Fytas, K., G.J. Georgopoulos, H. Soto, and V.P. Evangelou. Feasibility and cost of creating an iron phosphate coating on pyrrhotite to prevent oxidation. *Environmental Geology*, 28:61-69.
- Gan, S., and R.M. Amasino. Cytokinins in plant senescence: From spray and pray to clone and play. *BioEssays*, 18:557-565.
- Hadad, R.G., T.W. Pfeiffer, and C.G. Poneleit. Repeatability and heritability of divergent recombination frequencies in the Iowa stiff stalk synthetic (*Zea Mays* L.). *Theoretical and Applied Genetics*, 92:990-996.
- Hayati, R., D.B. Egli, and S.J. Crafts-Brandner. Independence of nitrogen supply and seed growth in soybean: Studies using an *in vitro* culture system. *Journal of Experimental Botany*, 47:33-40.
- Hong, Y., and A.G. Hunt. RNA polymerase activity catalyzed by a potyvirus-encoded RNA-dependent RNA polymerase. *Virology*, 226:146-151.
- Howell, J.H., M.S. Coyne, and P.L. Cornelius. Effect of sediment particle size and temperature on fecal bacteria mortality rates and the FC/FS ration. *Journal of Environmental Quality*, 25:1216-1220.
- Li, Q., J. Das Gupta, and A.G. Hunt. A plant poly(A) polymerase requires a novel RNA binding protein for activity. *Journal of Biological Chemistry*, 271:19831-19835.
- Liu, W., R.S. Torisky, K.P. McAllister, S. Avdiushko, D. Hildebrand, and G.B. Collins. Somatic embryo cycling: Evaluation of a novel transformation and assay system for seed-specific gene expression in soybean. *Plant Cell, Tissue and Organ Culture*, 47(1):33-42.
- Ma, Y.Z., C.T. MacKown, and D.A. Van Sanford. Differential effects of partial spikelet removal and defoliation on kernel growth and assimilate levels among wheat cultivars. *Field Crops Research*, 47:201-209.
- Ma, Y.Z., C.T. MacKown, and D.A. Van Sanford. Kernel growth of *in vitro* cultured wheat spikes of cultivars with divergent source-sink limitations. *Plant Science*, 119:135-148.
- MacKown, C.T., D.A. Van Sanford, and C.G. Rothwell. Nitrate uptake and assimilation and chlorate tolerance of wheat. *Crop Science*, 36:313-318.
- Miles, C.L., G.A. Lane, M.E. di Menna, I. Garthwaite, E.L. Piper, O.J.-P. Ball, G.C.M. Latch, J.M. Allen, M.B. Hunt, L.P. Bush, F.K. Min, I. Fletcher, and P.S. Harris. High levels of ergonovine and lysergic acid amide in toxic *Achnatherum inebrians* accompany infection by an *Acremonium*-like endophytic fungus. *Journal of Agriculture and Food Chemistry*, 44:1285-1290.
- Murphy, J.F., P.G. Klein, A.G. Hunt, and J.G. Shaw. Replacement of the tyrosine residue that links a potyviral VPg to the viral RNA is lethal. *Virology*, 220:535-538.
- Nadler, A., E. Perfect, and B.D. Kay. Effect of polyacrylamide application on the stability of dry and wet aggregates. *Soil Science Society of America Journal*, 60:555-561.
- Norfleet, M.L., and A.D. Karathanasis. Some physical and chemical factors contributing to fragipan strength in Kentucky soils. *Geoderma*, 71:289-301.
- Pearce, W.L., D.A. Van Sanford, and D.E. Hershman. Partial resistance to powdery mildew in soft red winter wheat. *Plant Diseases*, 80:1359-1362.
- Perfect, E., and N.B. McLaughlin. Soil management effects on planting and emergence of no-till corn. *Transactions, American Society of Agricultural Engineers (ASAE)*, 39(5):1611-1615.
- Perfect, E., N.B. McLaughlin, B.D. Kay, and G.C. Topp. An improved fractal equation for the soil water retention curve. *Water Resources Research*, 32:281-287.
- Pfeiffer, T.W., L.C. Harris, D.L. Pilcher, and C.R. Tutt. Registration of 'CF492' Soybean. *Crop Science*, 36:811-812.
- Pfeiffer, T.W., L.C. Harris, D.L. Pilcher, and J.M. Wood. Registration of 'CF461' soybean. *Crop Science*, 36:811.
- Pfeiffer, T.W. Choosing soybean varieties from yield trials: Multiple maturity groups and yield variability. *Journal of Production Agriculture*, 9:371-376.
- Sanderson, M.A., F. Agblevor, M. Collins, and D.K. Johnson. Compositional analysis of biomass feedstocks by near infrared reflectance spectroscopy. *Biomass and Bioenergy*, 11:365-370.
- Semalulu, O., and R.I. Barnhisel. Availability of P and K in restored prime farmland. *Proceedings, American Society for Surface Mining and Reclamation*, 13:442-450.
- Seta, A.K., and A.D. Karathanasis. Water dispersible colloids and factors influencing their dispersibility from soil aggregates. *Geoderma*, 74:255-266.
- Seta, A.K., and A.D. Karathanasis. Colloid-facilitated transport of metolachlor through intact soil columns. *Journal of Environmental Science and Health*, B31(15):949-968.
- Shen, P.-S., P.L. Cornelius, and R.L. Anderson. Planned unbalanced designs for estimation of quantitative genetic parameters. II. Nested matings. *Journal of Agricultural, Biological and Environmental Statistics*, 1:490-505.
- Shen, P.-S., P.L. Cornelius, and R.L. Anderson. Planned unbalanced designs for estimation of quantitative genetic parameters. I. Two-way matings. *Biometrics*, 52:56-70.
- TeKrony, D.M., L.J. Grabau, M. DeLacey, and M. Kane. Influence of early planted, early maturing soybeans on seed germination and *Phomopsis* seed infection. *Agronomy Journal*, 88:428-433.
- Thom, W.O., and R.L. Blevins. Conservation tillage and filter strips trap potential water contaminants. *Better Crops*, 80:12-14.
- Thomas, G.W., G.R. Haszler, and R.L. Blevins. The effects of organic matter and tillage on maximum compactability of soils using the Proctor test. *Soil Science*, 161:502-508.
- Thompson, J.A., and J.C. Bell. Color index for identifying hydric soil conditions in seasonally-saturated Mollisols. *Soil Science Society of America Journal*, 60:1979-1988.
- Torisky R., J. Fellers, and G.B. Collins. A focusing device for tissue transformation with the Dupont/BioRad PDS 1000 helium microprojectile system. *Plant Molecular Biology Reporter*, 14:124-133.
- Torisky, R.S., W. Liu, K.P. McAllister, S. Avdiushko, D. Hildebrand, and G.B. Collins. A mammalian desaturase gene lowers saturated fatty acid levels in transgenic soybean embryos. *Plant Cell, Tissue and Organ Culture*, 47:33-42.
- Vogeli-Lang, R., and G.J. Wagner. Relationship between Cd, GSH and Cd-binding peptides in leaves of tobacco seedlings. *Plant Science*, 114:11-18.
- Williams, D.W., A.J. Powell, P. Vincelli, and C.T. Dougherty. Dollar spot on bentgrass influenced by displacement of leaf surface moisture, nitrogen, and clipping removal. *Crop Science*, 36(5):1304-1309.
- Zhai, Q., and R.I. Barnhisel. A reclamation approach for mined prime farmland by adding organic wastes and lime. *Proceedings, American Society for Surface Mining and Reclamation*, 13:740-748.
- Zhai, Q., and R.I. Barnhisel. Crop and soil responses to sewage sludge applied to reclaimed prime farmland. *Proceedings, American Society for Surface Mining and Reclamation*, 13:813-820.
- Zhang, Zhiguo, and R.L. Blevins. Corn yield response to cover crops and N rates under long-term conventional and no-tillage management. *Journal of Sustainable Agriculture*, 8:61-72.
- Zhang, Y.L., and V.P. Evangelou. Influence of iron oxide forming conditions on pyrite oxidation. *Soil Science*, 161:852-864.

(In addition, members of the department published 125 abstracts.)

Animal Sciences

- Besong, S., J.A. Jackson, C.L. Hicks, and R.W. Hemken. Effects of supplemental liquid yeast product on feed intake, ruminal profiles, yield, composition, and organoleptic characteristics of milk from lactating Holstein cow. *Journal of Dairy Science*, 79:1654-1658.
- Boatright, W.L., and A.D. Crum. Non-polar-volatile lipids from soy protein isolates and hexane-defatted flakes. *Journal of the American Oil Chemists' Society*, 74:461.
- Bucholtz, D.C., N.M. Vidwans, C.G. Herbosa, K.K. Schillo, and D.L. Foster. Metabolic interfaces between growth and reproduction. V. Pulsatile LH secretion is dependent upon glucose availability. *Endocrinology*, 137:601-607.
- Carter, S.D., G.L. Cromwell, T.R. Combs, G. Colombo, and P. Fanti. The determination of serum concentrations of osteocalcin in growing pigs and its relationship to standard indices of bone integrity. *Journal of Animal Science*, 74:2719-2729.
- Cervantes, M.R., G.L. Cromwell, and T.S. Stahly. Synthetic amino acid supplementation in the low protein grain sorghum-soybean meal diet for pigs. *Cuban Journal of Agricultural Science*, 29:197-208.
- Chalupa, W., B. Vecchiarelli, D.T. Galligan, J.D. Ferguson, L.S. Baird, R.W. Hemken, R.J. Harmon, C.G. Soderholm, D.E. Otterby, R.J. Annexstad, J.G. Linn, W.P. Hansen, F.R. Ehle, D.L. Palmquist, and R.G. Eggert. Responses of dairy cows supplemented with somatotropin during weeks 5 through 43 of lactation. *Journal of Dairy Science*, 79:800-812.
- Cromwell, Gary L. Low protein, amino acid supplemented diets for growing-finishing pigs. pp. 13-26. Proceedings, 16th Agricultural Feed Ingredient Conference, Quincy, IL.
- Cromwell, Gary L. Using nutritional practices to reduce nutrient load and control odor. pp. 107-126. Proceedings, Pork Profitability Summit. National Pork Producers Council, Des Moines, IA.
- Cromwell, Gary L. The effects of phytase on phosphorus utilization in swine. pp. 18. BASF Technical Symposium, Des Moines, IA.
- Cromwell, Gary L. Optimizing phosphorus levels in growing-finishing diets for swine. pp. 13. American Association of Swine Practitioners, Nashville, TN.
- Cromwell, Gary L. Trace elements and animal production. 9th International Symposium on Trace Elements in Man and Animals, Banff, Alberta, Canada.
- Cromwell, Gary L. Swine nutrition today and in the next century. pp. 14. Seminario de Actualizacion en Nutricion Porcina. Irapuato, Guanajuato, Mexico.
- Cromwell, G.L., G.W. Davis, W.E. Morgan Morrow, R.A. Primo, D.W. Rozenboom, M.D. Sims, E.P. Stanisiewski, and C.H. Ho. Efficacy of the antimicrobial compound U-82, 127 as a growth promoter for growing-finishing pigs. *Journal of Animal Science*, 74:1284-1287.
- Du, Z., R.W. Hemken, and R.J. Harmon. Copper metabolism of Holstein and Jersey cows and heifers fed diets high in cupric sulfate or copper proteinate. *Journal of Dairy Science*, 79:1873-1880.
- Du, Z., R.W. Hemken, J.A. Jackson, and D.S. Trammel. Utilization of copper in copper proteinate, copper lysine, and cupric sulfate using the rat as an experimental model. *Journal of Animal Science*, 74:1657-1663.
- Harmon, R.J. Control of bovine mastitis. pp. 5-17. Proceedings, Alltech do Brasil Lecture Tour, Curitiba, Brazil (Portuguese).
- Harmon, R.J. Control of contagious mastitis. pp. 11-19. Proceedings, National Mastitis Council, Madison, WI.
- Harmon, R.J., and J.K. Reneau. Factors affecting somatic cell counts in milk. pp.18-27. Proceedings, Alltech do Brasil Lecture Tour, Curitiba, Brazil (Portuguese).
- Harmon, R.J., and P.M. Torre. Role of micronutrients in mastitis. pp. 28-39. Proceedings, Alltech do Brasil Lecture Tour, Curitiba, Brazil (Portuguese).
- Harper, A.F., M.D. Lindemann, and E.T. Kornegay. Fetal survival and conceptus development after 42 days of gestation in gilts and sows in response to folic acid supplementation. *Canadian Journal of Animal Science*, 76:157-160.
- Hussein, A.S., A.H. Cantor, A.J. Pescatore, and T.H. Johnson. Effect of dietary protein and energy levels on pellet development. *Poultry Science*, 75:973-978.
- Jackson, J.A., D.R. Varney, R.J. Petroski, R.G. Powell, L.P. Bush, M.R. Siegel, R.W. Hemken, and P.M. Zavos. Physiological responses of rats fed loline and ergot alkaloids from endophyte-infected tall fescue. *Drug and Chemical Toxicology*, 19:85-96.
- Krehbiel, C.R., R.A. Britton, D.L. Harmon, J.P. Peters, R.A. Stock, and H.E. Grotjan. Effects of varying levels of duodenal or midjejunal glucose and 2-deoxyglucose infusion on small intestinal disappearance and net portal glucose flux in steers. *Journal of American Science*, 74:693-700.
- Lindemann, Merlin. Chromium: Pro's/con's, differences in sources, insulin sensitivity, and sow work to date. pp. 1-11. Proceedings, 12th Annual Carolina Swine Nutrition Conference.
- Lindemann, Merlin. Current understanding of chromium use in swine diets. pp. 223-234. Proceedings, 57th Annual Minnesota Nutrition Conference.
- Lindemann, Merlin. Organic chromium—the missing link in farm animal nutrition? pp. 299-316. Proceedings, 12th Annual Symposium on Biotechnology in the Feed Industry.
- Liu, G., and Xiong, Y.L. Contributions of lipid and protein oxidation to rheological differences between chicken white and red muscle myofibrillar proteins. *Journal of Agricultural and Food Chemistry*, 44:779-784.
- Liu, G., and Xiong, Y.L. Storage stability of antioxidant-washed myofibrils from chicken white and red muscle. *Journal of Food Science*, 61:890-894.
- Lou, J., K.A. Dawson, and H.J. Strobel. Role of phosphorylytic cleavage in cellobiose and cellodextrin metabolism by the ruminal bacterium *Prevotella ruminicola*. *Applied Environmental Microbiology*, 62:1770-1773.
- Shantha, N.C., W.G. Moody, and Z. Tabeidi. Conjugated linoleic acid concentration in semimembranosus muscle of grass- and grain-fed and zeranol-implanted beef cattle. *Journal of Muscle Foods*, 8:105-110.
- Silvia, W.J., L.A. Edgerton, and M.A. Kaminski. Changes in uterine secretion of prostaglandin F₂ alpha in response to oxytocin during the estrous cycle, early pregnancy and estrogen-induced pseudopregnancy in swine. *Biological Reproduction*, 55:657-662.
- Srinivasan, S., and Xiong, Y.L. Gelation of beef heart surimi as affected by antioxidants. *Journal of Food Science*, 61:707-711.
- Srinivasan, S., and Xiong, Y.L. Sodium chloride mediated lipid oxidation in beef heart surimi-like material. *Journal of Agricultural and Food Chemistry*, 44:1697-1703.
- Swinkels, J.W.G.M., E.T. Kornegay, W. Zhou, M.D. Lindemann, K.E. Webb, Jr., and M.W.A. Verstegen. Effectiveness of a zinc amino acid chelate and zinc sulfate in restoring serum and soft tissue zinc concentrations when fed to zinc-depleted pigs. *Journal of Animal Science*, 74: 2420-2430.
- Torre, P.M., R.J. Harmon, R.W. Hemken, T.W. Clark, D.S. Trammel, and B.A. Smoth. Mild dietary copper insufficiency depresses blood neutrophil function in dairy cattle. *Journal of Nutritional Immunology*, 4(3):3-24.
- Walker, J.A., and D.L. Harmon. Technical note: A simple, rapid assay for beta-amylase in bovine pancreatic juice. *Journal of Animal Science*, 74:658-662.
- Xiong, Y.L. Impacts of oxidation on muscle protein functionality. 49:79-86. Proceedings, Reciprocal Meat Conference.
- Yi, Z., E.T. Kornegay, V. Ravindran, M.D. Lindemann, and J.H. Wilson. Effectiveness of Natuphos phytase in improving the bioavailabilities of phosphorus and other nutrients in soybean meal-based semipurified diets for young pigs. *Journal of Animal Science*, 74:1601-1611.

(In addition, members of the department published 56 abstracts.)

Biosystems and Agricultural Engineering

- Bader, M.J., L.G. Wells, and L.R. Walton. Monitoring the dynamics of scaled vehicles using a sonic digitizer. *Transactions, ASAE*, 39(2):435-441.
- Bucklin, R.A., M. Molenda, T.C. Bridges, and I.J. Ross. Slip-stick frictional behavior of wheat on galvanized steel. *Transactions, ASAE*, 39(2):649-653.

- Chao, K., and R.S. Gates. Design of switching control systems for ventilated greenhouses. *Transactions, ASAE*, 39(4):1513-1523.
- Colliver, D.G., and M.W. Liddament. Effects of dry-bulb and humidity design setpoints on theoretical energy used in ventilation in future buildings—the role of the services engineer. 1:79-87. London: CIBSE/ASHRAE.
- Desmond, E.D., A.D. Ward, N.R. Fausey, and S.R. Workman. Comparison of daily water table depth prediction by four simulation models. *Transactions, ASAE*, 39(1):111-118.
- Duncan, G.A., L.R. Walton, J.H. Casada, L.D. Swetnam, and B. Tapp. Development and evaluation of a cable-hoist system for housing burley tobacco. *Applied Engineering in Agriculture*, 12(4):411-416.
- Edwards, D.R., J.F. Murdoch, T.C. Daniel, and P.A. Moore, Jr. Quality of runoff from four northwest Arkansas pasture fields treated with organic and inorganic fertilizer. *Transactions, ASAE*, 39(5):1689-1696.
- Edwards, D.R., T.C. Daniel, and P.A. Moore, Jr. Vegetative filter strip design for grassed areas treated with animal manures. *Applied Engineering in Agriculture*, 12(1):31-38.
- Edwards, D.R., C.T. Haan, J.F. Murdoch, A.N. Sharpley, T.C. Daniel, and P.A. Moore, Jr. Application of simplified phosphorus transport models to pasture fields in northwest Arkansas. *Transactions, ASAE*, 39(2):489-496.
- Edwards, D.R., P.A. Moore, Jr., T.C. Daniel, and P. Srivastava. Poultry litter-treated length effects on quality of runoff from fescue plots treated with poultry litter. *Transactions, ASAE*, 39(1):105-110.
- Gates, R.S., D.G. Overhults, and S.H. Zhang. Minimum ventilation for modern broiler facilities. *Transactions, ASAE*, 39(3):1135-1144.
- Hardin, B.O., R.A. Bucklin, and I.J. Ross. 1995. Shear-beam analysis for seismic response of metal wheat bins. *Transactions, ASAE*, 39(2).
- Malone, R., R. Warner, and M. Byers. Runoff losses of surface-applied metribuzin as influenced by yard waste compost amendments, no-tillage, and conventional-tillage. *Bulletin of Environmental Contamination and Toxicology*, 57:536-543.
- Malone, R., R. Warner, and M. Byers. Subsurface losses of surface-applied metribuzin as influenced by yard waste compost amendments, no-tillage and conventional-tillage. *Bulletin of Environmental Contamination and Toxicology*, 57:751-758.
- Malone, R., R. Warner, M. Byers, D. Hilborn, and D. Gere. Extraction of metribuzin from soil using supercritical CO₂ (SFE). *Bulletin of Environmental Contamination and Toxicology*, 58:46-52.
- McKnight, R.H., C. Donnelly, H.P. Cole, I.J. Ross, and L.R. Piercy. Collaborating with a medical school to assess occupational health and safety content in an engineering curriculum. *Journal of Engineering Education*, October.
- Molenda, M., J. Horabik, and I.J. Ross. 1995. Wear-in effects on loads and flow in a smooth-walled bin. *Transactions, ASAE*, 39(1):225-231.
- Molenda, M., J. Horabik, and I.J. Ross. 1995. Effect of filling methods on load distribution in model grain bins. *Transactions, ASAE*, 39(1):219-224.
- Molenda, M., J. Horabik, and I.J. Ross. 1995. Reduction of wall friction in initial period of grain silo operation. pp. 165-172. *Proceedings, 10th International Conference on Reinforced and Posttensioned Concrete Silos and Tanks.*
- Nokes, S.E., F.M. Landa, and J.D. Hanson. Validation of the root zone water quality model (ver 2.1) from a crop growth perspective. *Transactions, ASAE*, 39(3):1177-1184.
- Payne, F.A., R.S. Gates, and B.F. Fannin. Measurement precision for a fiber optic diffuse reflectance sensor. *Transactions, ASAE*, 39(6):2193-2198.
- Schwab, C.V., I.J. Ross, G.M. White, and D.G. Colliver. Wheat loads and vertical pressure distribution in a full-scale bin: Part II - Detention. *Transactions, ASAE*, 39(3):1145-1149.
- Shearer, S.A., J.A. Thomasson, and S.G. McNeill. Filter selection method for NIR sensing of plant and soil materials. *Transactions, ASAE*, 39(3):1209-1214.
- Srivastava, P., D.R. Edwards, T.C. Daniel, P.A. Moore, Jr., T.A. Costello, and D.J. Nichols. Vegetative filter strip performance with varying pollutant source and filter strip lengths. *Transactions, ASAE*, 39(6):2231-2239.
- Srivastava, P., T.A. Costello, and D.R. Edwards. A direct, approximate solution to the modified green-ampt infiltration equation. *Transactions, ASAE*, 39(4):1311-1313.
- Stombaugh, D.P. and S.E. Nokes. Development of a biologically-based aerobic composting simulation model. *Transactions, ASAE*, 39(1):239-250.
- Turner, L.W., G.L. Cromwell, T.C. Bridges, S. Carter, and R.S. Gates. Ammonia (NH₃) emission from swine waste as influenced by diet manipulation. pp. 453-458. *Proceedings, 1st International Conference on Air Pollution from Agricultural Operations*. Ames, IA: Midwest Plan Service, Iowa State University.
- Wu, Q.J., A.D. Ward, and S.R. Workman. Using GIS in simulation of nitrate leaching from heterogeneous unsaturated soils. *Journal of Environmental Quality*, 25(3):526-534.
- Zhuang, H., F.A. Payne, S. Bastin, B.E. Langlois, and M.M. Barth. Ozone water treatments for preserving quality of packaged, fresh-cut broccoli under refrigeration. pp. 267-276. *Proceedings, Conference on New Developments in Refrigeration for Food Safety and Quality, ASAE.*
- (In addition, the department published eight abstracts.)
- ### Entomology
- Baker, T.C., and K.F. Haynes. Pheromone-mediated optomotor anemotaxis and altitude control exhibited by male oriental fruit moths in the field. *Physiological Entomology*, 21:20-32.
- Baur, M.E., and K.V. Yeargan. 1995. Host selection and larval interactions among three primary parasitoids of *Plathypena scabra* (Lep.: Noctuidae). *Entomophaga*, 40:357-366.
- Baur, M.E., and K.V. Yeargan. Movement and response to semiochemicals by parasitoids of *Plathypena scabra* (Lepidoptera: Noctuidae). *Journal of the Kansas Entomological Society*, 69:122-132.
- Baur, M.E., K.V. Yeargan, and P.L. Cornelius. Allocation of searching time within the soybean canopy by parasitoids attacking the green cloverworm. *Journal of Insect Behavior*, 9:223-235.
- Brown, G.C. Entomology and pest management. Book Review. *Journal of Economic Entomology*, 89:1669.
- Brown, G.C. Forecasting biological control effectiveness: A general case applied to aphid management. pp. 223-233. G. Bourgeois, and M.O. Guibord, eds. *Proceedings, International Symposium on Agricultural Pest Forecasting and Monitoring*, Quebec, Canada.
- Clark, D.C., S.J. DeBano, and A.J. Moore. The influence of environmental quality on sexual selection in *Nauphoeta cinerea* (Dictyoptera: Blaberidae). *Behavioral Ecology*, 8:46-53.
- Cui, L., and B.A. Webb. Isolation and characterization of a member of the cysteine-rich gene family from *Campoletis sonorensis* polydnavirus. *Journal of General Virology*, 77:797-809.
- Dangerfield, P.C., J.B. Whitfield, M.J. Sharkey, D.H. Janzen, and I. Mercado. *Hansonia*, a new genus of cardiochiline Braconidae (Hymenoptera) from Costa Rica, with notes on its biology. *Proceedings, Entomological Society of Washington*, 98:592-596.
- Dong, K., D. Zhang, and D.L. Dahlman. Down-regulation of juvenile hormone esterase and arylphorin production in *Heliothis virescens* larvae parasitized by *Microplitis croceipes*. *Archives of Insect Biochemistry and Physiology*, 32:237-248.
- Haynes, K.F., K.V. Yeargan, J.G. Millar, and B.B. Chastain. Identification of the sex pheromone of *Tetanolia mynesalis* (Lepidoptera: Noctuidae), a prey species of bolas spider, *Mastophora hutchinsoni*. *Journal of Chemical Ecology*, 22:75-89.
- Johnson, D.W., and D.E. Hershman. Survey of cereal aphids in Kentucky wheat fields: Common species and distribution. *Transactions, Kentucky Academy of Sciences*, 57:15-17.
- Kreiter, N., and D.H. Wise. Age-related changes in movement patterns in the fishing spider, *Dolomedes triton* (Araneae, Pisauridae). *Journal of Arachnology*, 24:24-33.

- Loughrin, J.H., D.A. Potter, T.R. Hamilton-Kemp, and M.E. Byers. Role of feeding-induced plant volatiles in aggregative behavior of the Japanese beetle (*Coleoptera: Scarabaeidae*). *Environmental Entomology*, 25:1188-1191.
- Loughrin, J.H., D.A. Potter, T.R. Hamilton-Kemp, and M.E. Byers. Volatile compounds from crabapple (*Malus* spp.) cultivars differing in susceptibility to the Japanese beetle (*Popillia japonica* Newman). *Journal of Chemical Ecology*, 22:1295-1305.
- Luckhart, S., and B.A. Webb. Interaction of a wasp ovarian protein and polydnavirus in host immune suppression. *Developmental and Comparative Immunology*, 20:1-21.
- Monroy, A.M., T.W. Scott, and B.A. Webb. Evaluation of Reverse Transcriptase Polymerase Chain Reaction for the detection of eastern equine encephalomyelitis virus during vector surveillance. *Journal of Medical Entomology*, 33:449-457.
- Potter, D.A., J.H. Loughrin, W.J. Rowe II, and T.R. Hamilton-Kemp. Why do Japanese beetles defoliate trees from the top down? *Entomologia Experimentalis et Applicata*, 80:209-212.
- Potter, D.A., A.J. Powell, P.G. Spicer, and D.W. Williams. Cultural practices affect root-feeding white grubs (*Coleoptera: Scarabaeidae*) in turfgrass. *Journal of Economic Entomology*, 89:156-164.
- Presley, S.M., F.W. Knapp, J.A. Boling, and J.G. Burg. Effects of the horn fly (*Diptera: Muscidae*) on physiological and nutritional responses of beef steers: Continuous fly population levels. *Journal of Economic Entomology*, 89:138-143.
- Rieske, L.K., and F.B. Peairs. Gustatory and olfactory responses of *Cylindrocopturus adspersus* (LeConte) to various plant species in the laboratory. *Southwestern Entomologist*, 21:255-264.
- Rieske, L.K., and K.F. Raffa. Bionomics and host range of the introduced basswood thrips (*Thysanoptera: Thripidae*). *Annals of the Entomological Society of America*, 89:75-80.
- Rowe, W.J. II, and D.A. Potter. Vertical stratification of feeding by Japanese beetles within linden tree canopies: Selective foraging or height per se? *Oecologia*, 108:459-466.
- Sharkey, M.J. The Agathidinae (Hymenoptera: Braconidae) of Japan. *Bulletin of the National Institute of Agro-Environmental Sciences*, 13:1-100.
- Sharkey, M.J. Cladistics of the ichneutinae revisited. *Journal of Natural History*, 30:1577-1578.
- Soldevila, A.I., and B.A. Webb. Expression of polydnavirus genes under polydnavirus promoter regulation in insect larvae infected with baculovirus recombinants. *Journal of General Virology*, 77:1379-1388.
- Wagner, J.D., and D.H. Wise. Cannibalism regulates densities of young wolf spiders: Evidence from field and laboratory experiments. *Ecology*, 77:639-652.
- Webb, B.A., and S. Luckhart. Factors mediating short- and long-term immune suppression in a parasitized insect. *Journal of Insect Physiology*, 42:33-40.
- Yeagan, K.V., and W.E. Barney. Photoperiodic induction of reproductive diapause in the predators *Nabis americanoferus* and *Nabis roseipennis* (Heteroptera: Nabidae). *Annals of the Entomological Society of America*, 89:70-74.
- Yeagan, K.V., and L.W. Quate. Juvenile bolas spiders attract psychodid flies. *Oecologia*, 106:266-271.
- (In addition, members of the department published 27 abstracts.)

Forestry

- Adams, J.P., M.J. Lacki, and M.D. Baker. Response of herpetofauna to silvicultural prescriptions in the Daniel Boone National Forest, Kentucky. *Proceedings, Southeastern Association of Fish and Wildlife Agencies*, Vol. 50.
- Arthaud, G.J., and M.H. Pelkki. A comparison of dynamic programming and A* in optimal forest stand management. *Forest Science*, 42:498-503.
- Arthaud, G.J., and M.H. Pelkki. The impact of production types and prices on optimal stand management. pp. 114-12. J. Caulfield and S. Bullard, eds. *Proceedings, 1995 Southern Forest Economists Workshop*, New Orleans. Mississippi State University Press.

- Gracey, E., and M.H. Pelkki. Comparing woodland owner association members to the nonindustrial private landowner population. pp. 403-410. M. Baughman, ed. *Proceedings, Symposium on Nonindustrial Private Forests*, Washington, D.C.
- Lacki, M.J. The role of research in conserving bats in managed forests. pp. 39-48. Work Paper 23. IN: R.M.R. Barclay and R.M. Brigham, eds. *Bats and Forests Symposium*. Res. Br. B.C. Min. For., Victoria, B.C.
- Pelkki, M.H. Adding a temporal dimension to stand-level forest databases. *The Compiler*, 14:14-23.
- Pelkki, M.H. Exploring spatial confidence in a raster-based GIS. *Proceedings, Spatial Accuracy Assessment in Natural Resources and Environmental Science: 2nd International Symposium*. USDA RM-GTR-277:336.342.
- Pelkki, M.H. Forest and e-mail use: A simple survey. *The Compiler*, 14:40-45.
- Pelkki, M.H. GIS and database management systems: Managing spatial and aspatial data. *The Compiler*, 14:26-33.
- Pelkki, M.H. Optimal timber management for shortleaf pine in the central United States. pp. 139-147. J. Caulfield and S. Bullard, eds. *Proceedings, 1995 Southern Forest Economists Workshop*, New Orleans. Mississippi State University Press.
- Pelkki, M.H. What does "relational" mean in a relational database management system? *The Compiler*, 14:38-40.
- Pelkki, M.H., J.M. Ringe, D.L. Brown, and D.H. Graves. Woody plant biodiversity on an abandoned coal washing pond. *International Journal of Surface Mining, Reclamation and Environment*, 10:161-166.
- Pelkki, M.H., J.M. Ringe, and D.H. Graves. Evaluation of shredded rubber tires as a surface mine spoil amendment. *International Journal of Surface Mining, Reclamation and Environment*, 10:55-59.
- Stringer, J.W., and D.H. Graves. Tree shelters overcome initial establishment problems for containerized seedlings of *Pawlonia tomentosa* over a wide range of planting conditions. pp. 74. *Proceedings, Tree Shelter Conference*. USDA Forest Service General Technical Report NE-221.
- (In addition, members of the department published 14 abstracts.)

Horticulture and Landscape Architecture

- Archbold, D.D. Evidence of annual variation in nitrogen assimilation by strawberry roots. *Advances in Strawberry Research*. 15:21-24.
- Archbold, D.D. Drought resistance in strawberry: What is the potential for improvement? pp. 127-132. M.P. Pritts et al., eds. *Proceedings, 4th North American Strawberry Conference*. Gainesville: University of Florida Press.
- Archbold, D.D., T.R. Hamilton-Kemp, B.E. Langlois, and M.M. Barth. Use of natural volatile compounds for controlling strawberry fruit quality. pp. 237-243. *Proceedings, Conference on New Developments in Refrigeration for Food Safety and Quality*. ASAE.
- Autio, W.R., R.A. Hayden, W.C. Micke, and G.R. Brown. Rootstock affects ripening, color, and shape of 'Starkspur Supreme Delicious' apples in the 1984 NC-140 Cooperative Planting.
- Crankshaw, N. Street trees in historic downtowns: A case study of Danville, Kentucky. *Association of Preservation Technology Bulletin*, 27(4):36-42.
- Galloway, B.A., and L.A. Weston. Influence of cover crop and herbicide treatment on weed control and yield in no-till sweet corn and pumpkins. *Weed Technology*, 10:341-346.
- Hamilton-Kemp, T.R., D.D. Archbold, J.H. Loughrin, R.W. Collins, and M.E. Byers. Metabolism of natural volatile compounds by strawberry fruit. *Journal of Agricultural and Food Chemistry*, 44:2802-2805.
- Hoffman, M.L., J. Buxton, and L.A. Weston. Using subirrigation to maintain soil moisture content in greenhouse experiments. *Weed Science*, 44: 397-401.
- Hoffman, M.L., L.A. Weston, J.C. Snyder, and E.E. Regnier. Separating the effects of sorghum (*Sorghum bicolor*) and rye (*Secale cereale*) root and shoot residues on weed development. *Weed Science*, 44:402-407.

- Hoffman, M.L., L.A. Weston, J.C. Snyder, and E.E. Regnier. Interference mechanisms between germinating seeds and between seedlings bioassays using cover crop and weed species. *Weed Science*, 44:579-584.
- Merkin, Z.R., and T.J. Nieman. Reinterpreting SMCRA: 'Permitting' phased postmining land use. pp. 766-780. W.L. Daniel, ed. *Proceedings, American Society for Surf Mining and Reclamation*.
- Nimbal, C.I., L.A. Weston, J. Pedersen, and C. Yerkes. Phytotoxicity and distribution of sorgoleone in grain sorghum germplasm. *Journal of Agricultural and Food Chemistry*, 44:1343-1347.
- Nimbal, C.I., C.N. Yerkes, L.A. Weston, and S.C. Weller. Herbicidal activity and site of action of the natural product sorgoleone. *Pesticide Biochemistry and Physiology*, 54:73-83.
- Nimbal, C.I., and L.A. Weston. Herbicidal activity and site of action of sorgoleone. *Proceedings, International Weed Congress, Copenhagen, Denmark*.
- Ying, Z., N. Janney, and R. Houtz. Organization and characterization of the ribulose-1,5-bisphosphate carboxylase/oxygenase large subunit N-methyltransferase gene in tobacco. *Plant Molecular Biology*, 32:663-671.
- Zhuang, H., T.R. Hamilton-Kemp, R.A. Andersen, and D.F. Hildebrand. The impact of alteration of polyunsaturated fatty acid levels on C6-aldehyde formation of *Arabidopsis* leaves. *Plant Physiology*, 111:805-812.
- (In addition, members of the department published 43 abstracts.)
- Nutrition and Food Science**
- Chen L.H., and S.M. Lin. Acetaminophen alters antioxidant defense enzyme activities in hepatocyte culture. *Biochemical Archives*, 12:33-42.
- Chen, L.H., C.Y. Huang, and Y. Osio. Large doses of ethanol and activities of liver enzymes which detoxify reactive oxygen species: Effects of antioxidants. *Biochemical Archives*, 12:95-104.
- Chen, L.H., S. Xi, and D.A. Cohen. Liver endogenous antioxidant defenses in mice fed AIN-76A diet and infected with murine AIDS. *Chemico-Biological Interactions*, 99:17-28.
- Chow, C.K. Vitamins and related components: Will increased intake provide greater protection against oxidative damage? *Nutrition*, 12:822-823.
- Gagliardi, A.R., B. Hennig, and D.C. Collins. Antiestrogens inhibit endothelial cell growth stimulated by angiogenic growth factors. *Anticancer Research*, 16:1101-1106.
- Gorman, L., L.P. Mercer, and B. Hennig. Growth requirements of endothelial cells in culture: Variations in serum and amino acid concentrations. *Nutrition*, 12:266-270.
- Haq, Akram-ul, Holly Bundrant, and L. Preston Mercer. Food intake is inversely correlated with central nervous system H1 receptor concentrations in male SD rats fed normal, low protein, low energy or poor quality protein diets. *Journal of Nutrition*, 126:3083-3089.
- Haq, Akram-ul, H. Bundrant, and L. Preston Mercer. Histamine H1 receptors found to exist in chicken brain. *Poultry Science*, 75:46.
- Hennig, B., M. Toborek, and C.J. McClain. Antiatherogenic properties of zinc: Implications in endothelial cell metabolism. *Nutrition*, 12:711-717.
- Hennig, B., M. Toborek, C.J. McClain, and J.N. Diana. Nutritional in vascular endothelial implications cell metabolism. *Journal of American College of Nutrition*, 15:345-358.
- Hennig, B., M. Toborek, S. Joshi-Brave, S.W. Barger, S. Barve, M.P. Mattson, and C.J. McClain. Linoleic acid activates NF-kB and induces NF-kB-dependent transcription in cultured endothelial cells. *American Journal of Clinical Nutrition*, 63:322-328.
- Hong, J.T., and H.P. Glauert. Comitogenicity of eicosanoids and the peroxisome proliferator ciprofibrate in cultured rat hepatocytes. *Journal of Cellular Physiology*, 169:309-319.
- Leung, L.K., and H.P. Glauert. Reduction of the concentrations of prostaglandins E₂ and F_{2g} and thromboxane B₂ in cultured rat hepatocytes treated with the peroxisome proliferator ciprofibrate. *Toxicology Letters*, 85:143-149.
- Li, Y., L.K. Leung, H.P. Glauert, and B.T. Spear. Treatment of rats with the peroxisome proliferator ciprofibrate results in increased liver NF-kB activity. *Carcinogenesis*, 17:2305-2309.
- Li, Y., L.K. Leung, B.T. Spear, and H.P. Glauert. Activation of hepatic NF-kB by phenobarbital in rats. *Biochemical, Biophysical Research Communication*, 229:982-989.
- Mercer, L. Preston, D.S. Kelley, H.M. Bundrant, Akram-ul Haq, and L.L. Humphries. Gender affects rat's central nervous system histaminergic responses to dietary manipulation. *Journal of Nutrition*, 126:3128-3135.
- Mercer, L. Preston, D.S. Kelley, and L.L. Humphries. Dietary induced anorexia: Involvement of the histaminergic system. *Journal of American College of Nutrition*, 15:223-230.
- Nilakantan, V., Y. Li, H.P. Glauert, and B.T. Spear. Increased liver-specific catalase activity in transgenic mice. *DNA and Cell Biology*, 15:625-630.
- Ramasamy, S., D.W. Lipke, G.A. Boissonneault, H. Guo, and B. Hennig. Oxidized lipid-mediated alterations in proteoglycan metabolism in cultured pulmonary endothelial cells. *Atherosclerosis*, 120:199-208.
- Rogusa, R.J., C.K. Chow, D.K. St. Clair, and J.D. Porter. Antioxidant enzymes: Extraocular limb and diaphragm muscle group-specific activity patterns in control and mdx mice. *Journal of Neurological Sciences*, 139:180-186.
- Slim, R.M., M. Toborek, B.A. Watkins, G.A. Boissonneault, and B. Hennig. Susceptibility to hepatic oxidative stress in rabbits fed different animal and plant fats. *Journal of American College of Nutrition*, 15:289-294.
- Tatum, V.L., and C.K. Chow. Antioxidant status and susceptibility of sickle erythrocytes to oxidative and osmotic stress. *Free Radical Research*, 25:133-139.
- Toborek, M., and B. Hennig. Dietary methionine imbalance, endothelial cell dysfunction and atherosclerosis. *Nutrition Research*, 16:1251-1266.
- Toborek, M., S.W. Barger, M.P. Mattson, S. Barve, C.J. McClain, and B. Hennig. Linoleic acid and TNF-alpha cross-amplify oxidative injury and dysfunction of endothelial cells. *Journal of Lipids Research*, 37:123-135.
- (In addition, members of the department published 35 abstracts.)
- Plant Pathology**
- Chung, K.-R., and C.L. Schardl. Sexual cycle and horizontal transmission of the grass symbiont, *Epichloe typhina*. *Mycological Research*, 101:295-301.
- Chung, K.-R., A. Leuchtmann, and C.L. Schardl. Inheritance of mitochondrial DNA and plasmids in the ascomycetous fungus, *Epichloe typhina*. *Genetics*, 142:259-265.
- Gao, S., G.H. Choi, L. Shain, and D.L. Nuss. Cloning and targeted disruption of *eng-1*, encoding the major *in vitro* extracellular endopolygalacturonase of the chestnut blight fungus. *Applied Environmental Microbiology*, 62:1984-1990.
- Gould, A.B., J.W. Hendrix, and R.S. Ferriss. Relationship of mycorrhizal activity to time following reclamation of surface mine land in western Kentucky. I. Propagule and spore population densities. *Canadian Journal of Botany*, 74:247-261.
- Hartman, J.R. Evaluation of multi-layer fruit bags for sooty blotch and flyspeck control, 1995. *Biological and Cultural Tests*, 11:38.
- Hartman, J.R., U.E. Jarlfors, W.M. Fountain, and R. Thomas. First report of bacterial leaf scorch caused by *Xylella fastidiosa* on sugar maple and sweetgum. *Plant Disease*, 80:1302.
- Huang, S., and S.A. Ghabrial. Organization and expression of the double-standard RNA genome of *Helminthosporium victoriae* 190S virus, a totivirus infecting a plant pathogenic filamentous fungus. *Proceedings, National Academy of Sciences USA*, 93:12541-12546.
- Nicolas, O., T.P. Pirone, and G.M. Hellmann. Construction and analysis of infectious transcripts from a resistance-breaking strain of tobacco vein mottling potyvirus. *Archives of Virology*, 141:1535-1552.
- Vincelli, P., J. Doney, and A.J. Powell. Management of spring dead spot of bermudagrass, 1994-95. *Biological and Cultural Tests*, 11:25.
- Vincelli, P., J. Doney, and A.J. Powell. Impact of mowing height and nitrogen fertility on brown patch of tall fescue, 1995. *Biological and Cultural Tests*, 11:34.

- Vincelli, P., J. Doney, and A.J. Powell. Evaluation of fungicides for control of brown patch and anthracnose, 1995. *Fungicide and Nematicide Tests*, 51:362.
- Vincelli, P., J. Doney, and A.J. Powell. Evaluation of fungicides for control of brown patch, 1995. *Fungicide and Nematicide Tests*, 51:365.
- Vincelli, P., J. Doney, and A.J. Powell. Evaluation of fungicides for control of dollar spot, 1995. *Fungicide and Nematicide Tests*, 51:363-4.
- Wang, R.Y., E.D. Ammar, D.W. Thornbury, J.J. Lopez-Moya, and T.P. Pirone. Loss of potyvirus transmissibility and helper component activity correlates with non-retention of virions in aphid stylets. *Journal of General Virology*, 77:861-867.
- Wang, R.Y., and T.P. Pirone. Potyvirus transmission is not increased by pre-acquisition fasting of aphids reared on artificial diet. *Journal of General Virology*, 77:3145-3148.
- Wang, R.Y., and T.P. Pirone. Mineral oil interferes with the retention of tobacco etch potyvirus in the stylets of *Myzus persicae*. *Phytopathology*, 86:820-823.
- Wu, X., and J.G. Shaw. Bidirectional uncoating of the genomic RNA of a helical virus. *Proceedings, National Academy of Sciences USA*, 93:2981-2984.

(In addition, members of the department published 33 abstracts.)

Rural Sociology

- Hennon, C., S. Gavazzi, P. Dyk, and P. McKenry. Parenting practices and interventions among marginalized families in Appalachia. *Family Relations*, 45:388-396.
- Hustedde, R.J. An evaluation of the national issues forum methodology for stimulating deliberation in rural Kentucky. *Journal of the Community Development Society*, 27:2.

Veterinary Science

- Adams, M.H., C.B. Baker, K.J. McDowell. Molecular cloning and sequencing of equine inhibin cDNA. *Animal Biotechnology*, 7:1-10.
- Albright-Fraser, D.G., R. Reid, V. Gerber, E. Bailey. Polymorphism of DRA among equids. *Immunogenetics*, 43: 315-317.
- Coogle, L., E. Bailey, R. Reid, and M. Russ. Equine dinucleotide repeat polymorphisms at loci LEX002, -003, -004, -005, -007, -008, -009, -010, -011, -013 and -014. *Animal Genetics*, 27:126-127.
- Coogle, L., R. Reid, E. Bailey. Equine dinucleotide repeat polymorphisms at loci LEX025-LEX033. *Animal Genetics*, 27:289-290.
- Coogle, L., R. Reid, E. Bailey. Equine dinucleotide repeat polymorphisms at loci LEX015-LEX024. *Animal Genetics*, 27: 217-218.
- Daly, J.M., A.C.K. Lai, M.M. Binns, T.M. Chambers, M. Barrandeguy, and J.A. Mumford. Recent worldwide antigenic and genetic evolution of equine H3N8 influenza A viruses. *Journal of General Virology*, 77:661-671.
- Fitzgerald, B.P. Effects of administration of N-methyl-D,L-aspartate (NMA) on gonadotropin secretion in untreated and steroid-treated ovariectomized mares during the breeding season and in intact and ovariectomized mares during anestrus. *Domestic Animal Endocrinology*, 13:211-218.
- Fraser, D.G., and E. Bailey. Demonstration of three DRB loci in a domestic horse family. *Immunogenetics*, 44:441-445.
- Granstrom, D.E. Equine protozoal myeloencephalitis. *Veterinary Forum*, 7:52-57.
- Granstrom, D.E. Equine protozoal myeloencephalitis. *New York Thoroughbred*, 1:59-60.
- Green, B.E., L.D. Foil, S.D. Hagius, and C.J. Issel. The stability of equine infectious anemia virus in *Aedes aegypti* (Diptera: Culicidae), *Stomoxys calcitrans* (Diptera: Muscidae), and *Tabanus fuscicostatus* (Diptera: Tabanidae) stored at -70C. *Journal of the American Mosquito Control Association*, 12:334-336.
- Grund, C.H., E.R. Lechman, N.A. Pezzulo, C.J. Issel, and R.C. Montelaro. Fine specificity of EIAV gp90-specific antibodies associated with protective and enhancing immune responses in experimentally infected and immunized ponies. *Journal of General Virology*, 77:435-442.

- Guruprasad, K., T. Blundell, S. Xie, J. Green, B. Szafranska, R.J. Nagel, K. McDowell, C.B. Baker, R.M. Roberts. Comparative modeling and analysis of amino acid substitutions suggests that the family of pregnancy-associated glycoproteins includes both active and inactive aspartic proteinases. *Journal of Protein Engineering*, 9:849-856.
- Harkins, J.D., G.D. Mundy, S.D. Stanley, W.E. Woods, and T. Tobin. Regulatory significance of procaine residues in plasma and urine samples: A preliminary communication. *Equine Veterinary Journal*, 28(2):121-125.
- Harkins, J.D., G.D. Mundy, S.D. Stanley, W.E. Woods, R.A. Sams, D.R. Richardson, and T. Tobin. Character and duration of physiological effects of intravenous isoxsuprine. *Equine Veterinary Journal*, 28(4):320-326.
- Harkins, J.D., G.D. Mundy, S.D. Stanley, W.E. Woods, W.A. Rees, K.N. Thompson, and T. Tobin. Determination of highest no-effect doses for the local anesthetic responses to cocaine, bupivacaine and benzocaine. *Equine Veterinary Journal*, 28:30-37.
- Hedges, J.F., U.B.R. Balasuriya, P.J. Timoney, W.H. McCollum, and N.J. MacLachlan. Genetic variation in open reading Frame 2 of field isolates and laboratory strains of equine arteritis virus. *Virus Research*, 42:41-52.
- Hong, C.B. Congenital polyalveolar lobe in three foals. *Journal of Comparative Pathology*, 115:85-88.
- Kennedy, M.A., E. Ramsey, V. Diderrich, L. Richman, G.P. Allen, and L.N. Potgieter. Encephalitis associated with a variant of equine herpesvirus-1 in a Thomson's gazelle. *Journal of Zoo and Wildlife Medicine*, 27:533-538.
- Langemeier, J.L., S.J. Cook, R.F. Cook, K.E. Rushlow, R.C. Montelaro, and C.J. Issel. Detection of equine infectious anemia viral RNA in plasma samples from recently infected and long-term inapparent carrier animals by PCR. *Journal of Clinical Microbiology*, 34(6):1481-1487.
- Lichtenstein, D., C.J. Issel, and R.C. Montelaro. Genomic quaspecies associated with the initiation of infection and disease in ponies experimentally infected with equine infectious anemia virus. *Journal of Virology*, 70(6):3346-3355.
- Lyons, E.T., and S.C. Tolliver. Transmission of helminths of lambs on a farm in central Kentucky in 1993 and 1994. *Transactions, Kentucky Academy of Sciences*, 57(2):92-100.
- Lyons, E.T., J.H. Drudge, S.C. Tolliver, and S.S. Collins. Some aspects of experimental infections of *Trichostrongylus axei* in domestic rabbits (*Oryctolagus cuniculus*). *Journal of Helminthology Society, Washington*, 63(2):275-277.
- Lyons, E.T., S.C. Tolliver, J.H. Drudge, S. Stamper, T.W. Swerczek, and D.E. Granstrom. Critical test evaluation (1977-1992) of drug efficacy against endoparasites featuring benzimidazole-resistant small strongyles (Population S) in Shetland ponies. *Veterinary Parasitology*, 66:67-73.
- Lyons, E.T., S.C. Tolliver, J.H. Drudge, S. Stamper, T.W. Swerczek, and D. E. Granstrom. A study (1977-1992) of population dynamics of endoparasites featuring benzimidazole-resistant small strongyles (Population S) in Shetland ponies. *Veterinary Parasitology*, 66:75-86.
- Lyons, E.T., T.W. Swerczek, S.C. Tolliver, and J.H. Drudge. Natural superinfection of *Parascaris equorum* in a stall-orphan horse foal. *Veterinary Parasitology*, 66:119-123.
- Lyons, E.T., W.V. Bernard, C.B. Hong, and T.W. Swerczek. The pathogenicity of small strongyle larvae in Thoroughbred yearlings on a Kentucky farm. *Veterinary Medicine*, 91:466-472.
- McDowell, K.J., T.V. Little, P.J. Timoney, M.H. Adams. Characterization of proteins in seminal plasma of stallions, geldings, and geldings supplemented with testosterone. *Research in Veterinary Science*, 61:33-37.
- Montelaro, R.C., C. Grund, M. Raabe, B. Woodson, R.F. Cook, S. Cook, and C.J. Issel. Characterization of protective and enhancing immune responses to equine infectious anemia virus resulting from experimental vaccines. *AIDS Research and Human Retroviruses*, 12(5):413-415.
- Petrites-Murphy, M.B., L.A. Robbins, J.M. Donahue, and B. Smith. Equine cryptococcal endometritis and placentitis with neonatal cryptococcal pneumonia. *Journal of Veterinary Diagnostic Investigation*, 8:383-386.

- Runge, V.M., J.W. Wells, N.M. Williams, J.F. Timoney, and C. Lee. The use of gadolinium-BOPTA on magnetic resonance imaging in brain infection. *Investigative Radiology*, 31:294-299.
- Sams, R.A., D.F. Gerkin, R.L. Detra, W.E. Woods, S.D. Stanley, T. Tobin, J.M. Yang, H-H Tai, A. Jeganathan, and D.S. Watt. Identification of metabolites of azaperone in horse urine. *Journal of Pharmacological Science*, 85:79-84.
- Stokes, A., D.G. Alber, R. Cameron, R. Marshall, G.P. Allen, and R.A. Killington. The production of a truncated form of baculovirus expressed EHV-1 glycoprotein C and its role in protection of C3H (H-2K) mice against virus challenge. *Virus Research*, 44:97-109.
- Timoney, J.F., and M. Guan. Characterization of murine monoclonal antibodies recognizing opsonic, mouse-protective, chaining and mucosally relevant epitopes on the M protein of *Streptococcus equi* subsp. *equi*. *Research in Veterinary Science*, 60:76-81.
- Timoney, P.J., and W.H. McCollum. Equine viral arteritis. *Equine Veterinary Education*, 8:97-100.
- Timoney, P.J. Contagious Equine Metritis. *Comparative Immunology & Microbiology in Infectious Diseases*, 19 (3):199-204.
- Timoney, P.J. Equine Influenza. *Comparative Immunology & Microbiology in Infectious Diseases*, 19(3):205-211.
- Wellington, J.E., G.P. Allen, A.A. Gooley, D.N. Love, N.H. Packer, J.X. Yan, and J.M. Whalley. The highly O-glycosylated glycoprotein gp2 of equine herpesvirus 1 is encoded by gene 71. *Journal of Virology*, 70:8195-8198.
- Williams, N., L.R. Harrison, and C.J. Gebhart. Proliferative enteropathy in a foal caused by *Lawsonia intracellularis*-like bacterium. *Journal of Veterinary Diagnostic Investigation*, 8:254-256. (In addition, the department published nine abstracts.)

Other Research in Nonrefereed Publications

Agronomy

- Barbayiannis, N., V.P. Evangelou, and V.Z. Keramidas. Potassium-ammonium-calcium quantity/intensity studies in the binary and ternary modes in two soils of micaceous mineralogy. *Proceedings, Soil Science Society of Greece, Patra, Greece*.
- Barnhisel, R.I., O. Semalulu, and S. Witt. Determination of success in direct vegetation of ash disposal areas. pp. 1-20. *Kentucky Coal Utilization Conference*. Lexington, KY.
- Barnhisel, R.I., M.J. Bitzer, J.H. Grove, and S.A. Shearer. Agronomic benefits of varying corn seed populations: A central Kentucky study. *Proceedings, 3rd International Conference, Site-specific Management for Agriculture Systems*. American Society of Agronomy.
- Bitzer, M.J., R.I. Barnhisel, and J.H. Grove. Varying corn populations according to depth of topsoil. pp. 1-2. *Proceedings, Information Agriculture Conference*, Urbana, IL.
- Bitzer, M., and W. Bruening. Effect of varieties, row width and nitrogen rates on kenaf yields. pp. 92-95. *Proceedings, 8th International Kenaf Association Conference*, Albuquerque, NM.
- Chappell, J. Plant biotechnology comes of age—again. *Molecular Breeding*, 1:1-6.
- Collins, M. Advances in harvesting and storing high quality alfalfa. *Kentucky Alfalfa Conference*, 16:15-22.
- Collins, M., J. Henning, and G. Lacefield. Principles and challenges of making baled silage. pp. 26-31. *Proceedings, Kentucky Forage and Grassland Conference*, Lexington and Princeton, KY.
- Collins, M. Value of improved hay storage. pp. 32-34. *Proceedings, Kentucky Forage and Grassland Conference*, Lexington and Princeton, KY.
- Coyne, M.S., R.L. Blevins, J.H. Grove, D.R. Edwards, A. Pescatore, and A. Cantor. Nonpoint source pollution by land applied poultry waste: Runoff, infiltration, and management. pp. 12-33. In: W.O. Thom, ed. *Water quality research and education phase II: Progress report, first year*. Senate Bill 271 Water Quality Project Committee, University of Kentucky, Lexington, KY.
- Culton, E., Y. T. Wang, and W.O. Thom. Wetlands, an inexpensive means for wastewater treatment. pp. 21-23. *Proceedings, Annual Symposium, Water Resources and Research Institute*.
- Dougherty, C.T., and N. L. Taylor. The role of clovers in Kentucky grasslands. pp. 21-23. *Proceedings, 14th Trifolium Conference*. Lexington, KY.
- Evangelou, V.P. Remediation of acid drainage through surface coating of iron sulfides. 129 pp. U.S. Geological Survey, Department of the Interior.
- Evangelou, V.P. Formulation enhanced transport of a soil applied herbicide. University of Kentucky, Water Resources and Research Institute.
- Fairchild, M.A., M.S. Coyne, J.H. Grove, and W.O. Thom. Water quality and denitrification in perched water tables above fragipans. pp. 59-60. *Proceedings, Annual Symposium, Kentucky Water Resources*, Lexington, KY.
- Fytas, K., D.H. Soto, and V.P. Evangelou. (In French) Enrobage de la pyrite par le phosphate ferrique. pp. 191-213. *Proceedings, Conference on Acid Mine Drainage, Universite du Quebec-Rouyn-Noranda*.
- Grove, J.H., C.S. Stoddard, A.T. Basheeruddin, and W.O. Thom. Nitrate leaching under corn is not well related to choice of conservation tillage system. pp. 47-48. *Proceedings, Southern Conservation Tillage Conference for Sustainable Agriculture*, Jackson, TN.
- Grove, J.H., M.J. Bitzer, R.I. Barnhisel, and W.W. Frye. Predicting corn productivity for eroded upland landscapes in Kentucky: Soil clay and organic carbon contents. pp. 7-9. *Proceedings, Information Agriculture Conference*, Urbana, IL.
- Grove, J.H., and W.O. Thom. What have we learned about the role of conservation tillage on the quantity of nitrate nitrogen leaving the root zone of well-drained corn soils in Kentucky? pp. 7-8. *Proceedings, Annual Symposium, Kentucky Water Resources*.
- Legg, P. D. Physical components of quality in dark tobacco lines with different level of nicotine. pp. 39-40. *Tobacco Chemists' Research Conference*.
- Palmer, G., and R. Pearce. Major variety selections. *Burley and Dark Tobacco Production Guide 1996*. Raleigh, NC: Rural Press USA.
- Palmer, G., and R. Pearce. Tank-mix combinations show advantages. *Burley and Dark Tobacco Production Guide 1996*. Raleigh, NC: Rural Press USA.
- Pearce, R., and G. Palmer. Selecting the right fertilizer for the float system. *Burley and Dark Tobacco Production Guide 1996*. Raleigh, NC: Rural Press USA.
- Pearce, R., and G. Palmer. Measuring fertility of float beds. *Burley and Dark Tobacco Production Guide 1996*. Raleigh, NC: Rural Press USA.
- Pearce, R., R. Phillips, and J. Zeleznik. No-till burley tobacco production. *Burley and Dark Tobacco Production Guide 1996*. Raleigh, NC: Rural Press USA.
- Pfeiffer, T.W., S.A. Ghabrial, and Q. Ren. Resistance to soybean mosaic virus (SMV) is valuable for late-planted soybean. *Grain Gleamings*, 4(2).
- Sanderson, M.A., F. Agblevor, M. Collins, and D.K. Johnson. Compositional analysis of biomass feedstocks by near infrared reflectance spectroscopy. pp. 37-42. In: *Liquid Fuels and Industrial Products from Renewable Resources*. *Proceedings, 3rd National Bioenergy Conference*, Nashville, TN.
- Seta, A.K., and V.P. Evangelou. Pyrite microencapsulation: Potential for abatement of acid mine drainage. *Proceedings, 13th American Society of Surface Mining and Reclamation*, Knoxville, TN.
- TeKrony, D.M. Accelerated aging conditions for hybrid corn seed. *Iowa Seed Science Newsletter*, 16:4-5.
- TeKrony, D. M. Seed germination and vigor tests: How should results be utilized? *Seed World*, 134:32-34.
- Thom, W.O., R.L. Blevins, and M.S. Coyne. Use of filters to reduce non-point pollution. pp. 8-11. *Proceedings, Southern Soil Fertility Conference*. Ardmore, OK: Samuel Roberts Noble Foundation, Inc.
- Thom, W.O., Y.T. Wang, and J. Dinger. Evaluation of constructed wetlands for animal waste management. pp. 59-60. *Proceedings, Nonpoint Source Symposium*, Lexington, KY.
- Wells, K.L., and C.T. Dougherty. Soil considerations in a grazing system. pp. 63-68. *Proceedings, 1st Kentucky Grazing Symposium*.

Zourarakis, D.P., G.M. Young, S.A. Coleman, and W.O. Thom. Biogas production in Kentucky: Non-agricultural BMP alternatives for nonpoint source pollution prevention. pp. 71-78. Proceedings, 7th National Bioenergy Conference, Nashville, TN.

Animal Sciences

Bauer, M.L., and D.L. Harmon. Small intestinal glucose transport in ruminants: Effect of adaptation to digesting starch. pp. 83-86. IN: *1996 Beef Research Report* (PR-395), Agricultural Experiment Station, College of Agriculture, University of Kentucky, Lexington, KY.

Bohnert, D.W., M.L. Bauer, D.L. Harmon, G.E. Mitchell, Jr., and B.T. Larson. Ruminant degradation of poultry by-product meal. pp. 65-68. IN: *1996 Beef Research Report* (PR-395), Agricultural Experiment Station, College of Agriculture, University of Kentucky, Lexington, KY.

Cromwell, G.L., S.D. Carter, M.D. Lindemann, and L.W. Turner. Reducing the excretion of nitrogen and phosphorus in growing and finishing pigs by dietary manipulation. Proceedings, International Pig Veterinary Society Congress. Bologna, Italy.

Harmon, D.L., K.A. Dawson, M.L. Bauer, K.R. McLeod, and B.T. Larson. Interactions of yeast culture and protein supplementation for growing steers fed corn silage. pp. 45-54. IN: *1996 Beef Research Report* (PR-395), Agricultural Experiment Station, College of Agriculture, University of Kentucky, Lexington, KY.

Larson, B.T., D.L. Harmon, and R.M. Berry. Ergot alkaloid binding at D2 dopamine receptors in pituitary cells. pp. 91-96. IN: *1996 Beef Research Report* (PR-395), Agricultural Experiment Station, College of Agriculture, University of Kentucky, Lexington, KY.

Rowell, P.P., B.T. Larson, and D.L. Harmon. Ergot alkaloid-induced dopamine release from rat striatal synaptosomes. pp. 97-100. IN: *1996 Beef Research Report* (PR-395), Agricultural Experiment Station, College of Agriculture, University of Kentucky, Lexington, KY.

Walker, J.A., and D.L. Harmon. Influence of ruminal or abomasal starch hydrolysate infusion on pancreatic exocrine secretion and blood glucose and insulin concentration in steers. pp. 73-82. IN: *1996 Beef Research Report* (PR-395), Agricultural Experiment Station, College of Agriculture, University of Kentucky, Lexington, KY.

Biosystems and Agricultural Engineering

Colliver, D.G., H. Zhang, R.S. Gates, and K.T. Priddy. Sequences of Extreme Temperature and Humidity for Design Conditions. 89 pp. with 450MB CD-ROM. ASHRAE Research Project 828-RP. Atlanta, GA: American Society of Heating, Refrigerating and Air-Conditioning Engineers.

Dinger, J.S., D.M. Keagy, S.A. Minns, J.L. Taraba, and J.H. Grove. Assessment of nitrate and pesticide impacts on bedrock aquifers in upland agricultural settings of the western coal field. Water Quality Research and Education Program: Senate Bill 271 - Phase II Annual Progress Report. W.O. Thom, ed. College of Agriculture, University of Kentucky, Lexington, KY.

Edwards, D.R., G.K. Felton, J.L. Taraba, L.W. Turner, and B.T. Larson. Runoff quality responses to cattle grazing strategy and grassed buffer zone length. Water Quality Research and Education Program: Senate Bill 271 - Phase II Annual Progress Report. W.O. Thom, ed. College of Agriculture, University of Kentucky, Lexington, KY.

Fulton, J.P., L.G. Wells, S.A. Shearer, and R.I. Barnhisel. Spatial variation of soil physical properties: A precursor to precision tillage. ASAE Paper No. 961002, International Meeting, Phoenix, AZ.

Shearer, S.A., R.I. Barnhisel, S.F. Higgins, G.A. Watkins, and C.J. Sobolik. Assessment for the potential of site-specific management of crop production in central Kentucky. ASAE Paper No. 961003, International Meeting, Phoenix, AZ.

Shearer, S.A., R.I. Barnhisel, and G.A. Watkins. Site-specific management of herbicides on crop lands. pp. 34-48. Water Quality Research and Education, Phase II Progress Report for Senate Bill 271.

Sobolik, C.J., S.A. Shearer, and R.I. Barnhisel. Correlation of yield and soil parameters for variable-rate seeding of corn. ASAE Paper No. 961004, International Meeting, Phoenix, AZ.

Sun, S., and W.E. Murphy. Experimental and analytical evaluation of a flat plate type geothermal ground heat exchanger. Final Report to the National Rural Electric Cooperative Association.

Taraba, J.L., J.S. Dinger, L.V.A. Sendlein, and G.K. Felton. Land use impacts on water quality from small karst agricultural watersheds in Kentucky. Proceedings, 41st Midwest Groundwater Conference, Lexington, KY.

Warner, R.C., A.N. Hootkany, C.D. Blanton, J.L. Taraba, M.E. Byers, and L. Jarrett. Subsurface leaching potential of animal waste holding as function of soil moisture and compaction. Proceedings, Kentucky Nonpoint Source Conference, Lexington, KY.

Entomology

Potter, D.A., K.F. Haynes, and A.J. Powell. Cultural control, risk assessment, and environmentally responsible management of white grubs and cutworms in turfgrass. United States Golf Association. Turfgrass and Environmental Research Summary.

Potter, M.F. Talking "termites" with customers. *Pest Control*, 64(4):42,44,46.

Williamson, R.C., and D.A. Potter. Does aerification promote black cutworm infestations? *Golf Course Management*, January 97:54-57.

Horticulture and Landscape Architecture

Collins, K.B., L.A. Weston, and R.E. McNeil. Use of sulfentrazone (F6285) for weed management in field-grown ornamentals. 41:79-82. Proceedings, Southern Nurserymen's Association Research Conference.

Dunwell, W.C., W. Wolfe, and J. Johnston. Winter survival of daylily cultivars in west Kentucky. 41:348-350. Proceedings, Southern Nurserymen's Association Research Conference.

Finneseth, C.L.H., D.R. Layne, and R.L. Geneve. Germination and seedling emergence in pawpaw [*Asimina trioloba* (L.) Dunal]. 41:61-64. Proceedings, Southern Nurserymen's Association Research Conference.

Ingram, D.L., B. Roach, and M. Witt. Kentucky horticulture survey and database. 41:201. Proceedings, Southern Nurserymen's Association Research Conference.

Marohnic, J.J., R.L. Geneve, and J.W. Buxton. Capillary mats modify media moisture content during mist propagation. 41:99-102. Proceedings, Southern Nurserymen's Association Research Conference.

McNeil, R.E., L.A. Weston, and R. Harmon. Winter annual weed suppression by spring applied preemergence herbicide carryover. 41:251-252. Proceedings, Southern Nurserymen's Association Research Conference.

McNeil, R.E., L.A. Weston, and R. Harmon. Preemergence herbicides for use in annual bedding plants. 41:253-257. Proceedings, Southern Nurserymen's Association Research Conference.

NC-140 Committee. Performance of the NC-140 Cooperative Apple Rootstock Planting: I. Survival, Tree Size, and Fruit Size.

NC-140 Committee. Performance of the NC-140 Cooperative Apple Rootstock Planting: II. A 10-year Summary of TCA, Yield and Yield Efficiency at 31 Sites.

NC-140 Committee. Horticultural Characteristics of 'Starkspur Supreme Delicious' Apple on 15 Rootstocks at 24 Sites.

Roach, B.R., and R.E. McNeil. Price Trends of Nursery Stock. 41: 16-19. Proceedings, Southern Nurserymen's Association Research Conference.

Rowell, B., R.T. Jones, and W. Nesmith. Performance of new bell pepper cultivars exposed to induced and natural bacterial leaf spot epidemics. pp. 78-79. Proceedings, National Pepper Conference.

Rowell, B., R.T. Jones, J.C. Snyder, and W. Nesmith. Performance of new bacterial spot resistant pepper cultivars in Kentucky. pp. 81-84. Proceedings, Georgia State Vegetable Growers Association Annual Meeting.

- Stafford, M., R.L. Geneve, and R.E. McNiel. Changes in root length and diameter in plants grown in copper-treated containers. 41:73-74. Proceedings, Southern Nurserymen's Association Research Conference.
- Weston, L.A., and R.E. McNiel. Special report: Kentucky weed control results for ornamentals, 1996. College of Agriculture, University of Kentucky, Lexington, KY. 88 pp.

Plant Pathology

- Hartman, J., R. Bessin, G. Brown, R. Jones, and J. Strang. La protection integree pour la gestion des maladies des pommes dans le Kentucky, 1992 a 1996. IVE Congres de la Societe Francaise de Phytopathologie.
- Vincelli, P. Protecting tall fescue from brown patch. *Lawn & Landscape*, April 96:68-75.
- Vincelli, P. Disease diagnosis: Brown patch of tall fescue. *Turf and Ornamental Service Tech*, July 96:12.

Rural Sociology

- Burmeister, L.L. Comments on D. Gale Johnson's integration of East Asian agriculture into the world trading community. *Economic and Regional Cooperation in Northeast Asia*. 6:70-71. Washington, DC: The Korea Economic Institute of America.
- Burmeister, L.L. 1995. U.S. agriculture in the aftermath of the Uruguay round: Challenges and responses. pp. 25-45. *Korean Rural Sociological Society International Symposium on Social Change in Rural Societies and Countermeasures under the WTO System*. Seoul, Korea: KRSS.
- Dyk, P., and J. Ford. Family Preservation and Family Support Services Evaluation Final Report. University of Kentucky College of Agriculture, 42 pp.
- Harris, R., and others. 1995. Families First: Empowering Kentucky's People and Communities. Research Report No. 273, Legislative Research Commission, Frankfort, KY.
- Swanson, L.E., and others. RUPRI White Paper on Rural Policy: Rural Development Provisions of the 1996 Farm Bill and the Reorganization of USDA. 30 pp. Columbia, MO: Rural Policy Research Institute.
- Swanson, L.E., and others. Assessment of the House of Representatives Rural Development Proposals. 15 pp. Columbia, MO: Rural Policy Research Institute.

Veterinary Science

- Allen G. Equine Rhinopneumonitis. Chapter 3.4.7, pp. 1-8. IN: *OIE Manual of Standards for Diagnostic Tests and Vaccines*. Lyon: OIE Press.
- Balauriya, U.B.R., P.J. Timoney, W.H. McCollum, P.V. Rossitto, J.F. Patton, and N.J. MacLachlan. Phylogenetic analysis of open reading frame 5 and identification of major neutralization sites in the GL envelope glycoprotein of equine arteritis virus. p. 188. Abstract, 15th Annual Meeting of the American Society for Virology, Ontario.
- Cordes, T., and C.J. Issel. Equine infectious anemia: A status report on its control, 1996. 19 pp. United States Department of Agriculture Animal and Plant Health Inspection Service Publication 91-55-032.
- Dwyer, R.M. 1996 AAEP Convention Roundup. *The Horse*, January, pp. 49-54.
- Dwyer, R.M. American College of Veterinary Internal Medicine Forum Report. *The Horse*, August, p. 11.
- Dwyer, R.M. Commentary. *Equine Disease Quarterly*, October.
- Dwyer, R.M. Disinfection of equine facilities. *Equine Disease Quarterly*, July.
- Dwyer, R.M. Equine rotavirus vaccine. *Standardbred Breeders Newsletter*, August.

- Dwyer, R.M. On the road again (article on equine transportation). *The Horse*, March, p. 70.
- Dwyer, R.M. *Rhodococcus equi*: A killer that lurks in the dust. *The Horse*, October, pp. 33-36.
- Dwyer, R.M. Rotavirus vaccine. *Equine Disease Quarterly*, April.
- Dwyer, R.M. VEE: The current outbreak. *Equine Disease Quarterly*, January.
- Dwyer, R.M., M. Donahue. Equine salmonellosis: Central Kentucky and U.S. *Equine Disease Quarterly*, October.
- Granstrom, D.E. Equine protozoal myeloencephalitis: Clinical diagnosis and treatment. pp. 140-141. Proceedings, Michigan Veterinary Conference.
- Granstrom, D.E. Equine protozoal myeloencephalitis: Laboratory diagnosis. pp. 133-135. Proceedings, Michigan Veterinary Conference.
- Issel, C.J., Cook, R.F., and Montelaro, R.C. Equine infectious anemia. pp. 295-323. IN: M. Studdert, ed. *Equine Viruses*. Elsevier Press.
- Issel, C.J., Lane D. Foil, and Tim Cordes. Equine infectious anemia: Protecting your horse. Videotape produced by the USDA in collaboration with Louisiana State University and the University of Kentucky. Running time: 28 minutes.
- Kheyar, A.K., S. Martin, G. St-Laurent, W.H. McCollum, P.J. Timoney, and D. Archambault. Immunological Reactivity of the Equine Arteritis Virus Recombinant Nucleocapsid and Membrane Proteins. p. 112. Abstract, 77th Conference of Research Workers in Animal Diseases, Chicago, IL.
- Marti, E., G. Szalai, D.F. Antczak, E. Bailey, H. Gerber, and S. Lazary. The equine major histocompatibility complex. pp. 245-267. IN: L. Shook and S.J. Lamont, eds. *The Major Histocompatibility Complex of Domestic Animals*. Boca Raton, FL: CRC Press.
- Monreal, L., A.J. Villatoro, H. Hooghuis, and P.J. Timoney. Outbreak of equine viral arteritis in Spain. p. 15. Abstract, 1st European Symposium on Equine Viral Arteritis, Utrecht, The Netherlands.
- Monreal, L., A.M. Angles, M. Monreal, J. Monasterio, and P.J. Timoney. Changes in haemostasis in horses affected by EVA. p. 28. Abstract, 1st European Symposium on Equine Viral Arteritis, Utrecht, The Netherlands.
- Powell, D.G. Epidemiology and Control of Equine Viral Disease. Chapter 27. pp. 327-245. IN: M.J. Studdert, ed. *Virus Infections of Equines*. Elsevier, Amsterdam.
- Rees, W.A., J.D. Harkins, W.E. Woods, R.A. Blouin, M. Lu, C. Fenger, R.E. Holland, T. Tobin, and T.M. Chambers. Chemotherapy for equine influenza: Overview of the pharmacology and side effects of amantadine and rimantadine in the horse. pp. 355-357. Proceedings, 11th International Conference of Racing Analysts and Veterinarians, Queensland, Australia.
- Rees, W.A., S. Kwiatkowski, S.D. Stanley, D.E. Granstrom, J-M Yang, C.G. Gairola, J.N. Diana, J. Glowczyk, W.E. Woods, and T. Tobin. Development and characterization of an ELISA for trans-3'-hydroxycytidine. pp. 149-162. IN: Bengston, Henshel, eds. *Environmental Toxicology and Risk Assessment: Biomarkers and Risk Assessment* (5th vol.), American Society Testing Materials, Philadelphia.
- Timoney, P.J., and W.H. McCollum. Current strategies for the prevention and control of equine viral arteritis. pp. 121-125. IN: *Vaccines & Control of Infectious Diseases: The Way Forward*. Proceedings, 14th International Symposium, World Association of Veterinary Microbiologists, Immunologists, and Specialists in Infectious Diseases, Edinburgh.
- Timoney, P.J., B. Klingeborn, and M.H. Lucas. A perspective on equine viral arteritis (infectious arteritis of horses). *Revue Scientifique et Technique* (International Office des Epizooties), 15(3):1203-1208.
- Timoney, P.J. Client Information Series(tm), Equine Viral Arteritis. *Equine Practice*, 18:31-34.
- Timoney, P.J. Re-emergence in Mexico of Venezuelan equine encephalomyelitis (VEE). *Equine Disease Quarterly*, 5(1):3-4.

Ph.D. Dissertations

(Unpublished)

Agricultural Economics

Griffin, Peter W. Investigating the conflict in agricultural policy between the Federal Crop Insurance and Disaster Assistance Programs and the Conservation Reserve Program.

Ngarambe, Octavien. Income inequality vs. economic growth in United States: Examining causal relations.

Agronomy

Fellers, John P. Potyvirus Nia-Proteinase derived resistance: Evaluation of resistance and Nia interaction with the potyvirus polymerase.

Iin, Handayani. Soil carbon and nitrogen pools and transformations after 25 years of no tillage and conventional tillage.

Prawito, Priyono. Pedogenesis on Karst toposequences of Kentucky.

Ren, Q. Breeding soybean for double-cropping: Soybean mosaic virus resistance and pubescence density effects.

Xu, Dongmei. Genetic engineering strategies for virus resistance: Coat protein mediated resistance in tobacco (*Nicotiana tabacum* L.).

Animal Sciences

Bauer, Marc. Nutritional regulation of small intestinal glucose absorption in ruminants.

Besong, Samuel. Influence of supplemental chromium picolinate on the concentrations of hepatic triglyceride and blood metabolites in dairy cattle.

Girard, Ivan. Characterization of stimulatory activities from *Saccharomyces cerevisiae* on the growth and activities of ruminal bacteria.

Simeone, Ann-Marie. Ammoniation as a detoxification procedure for endophyte-infected tall fescue.

Spring, Peter. Effects of mannanoligosaccharide of different cecal parameters and on cecal concentrations of enteric pathogens in poultry. (Joint degree program with Swiss Federal Institute of Technology, Zurich).

Biosystems and Agricultural Engineering

Chao, Kuanglin. Economic optimization of single stem rose production.

Malone, Robert. Field assessment and modeling surface and subsurface metribuzin transport under three treatments: Yard waste compost amendments, no-till and conventional-till.

Entomology

Beavers, G.M. Utilization of low altitude remote sensing imagery to detect larval habitats of tire-breeding mosquitoes.

Cui, L. Functional studies of the cysteine-rich gene family of the *Campoletis sonorensis* (Hymenoptera: Ichneumonidae) polydnavirus.

Williamson, R.C. Behavior and ecology of the black cutworm in golf course turf.

Horticulture and Landscape Architecture

Distabanjong, Karsedis. Developing a genetic transformation system in Eastern redbud.

Plant Pathology

Chung, Kuang-Ren. Genetics of the grass symbiont, *Epichloe typhina* (Pers.: Fr.) Tul.

Huang, Shaohua. Organization and expression of the dsRNA genome of a dsRNA totivirus infecting the plant pathogenic fungus *Helminthosporium victoriae*.

Xie, Chengsong. Comparison of tobacco mosaic virus (TMV) and sodium salicylate as inducers of defense compounds and resistance to TMV and *Peronospora tabacina* in tobacco.

Veterinary Science

Mitchell, Eddie E. Addressing the regio- and stereo-selectivity seen in P450A5 mutants with DHEA.

Fenger, Clara K. Application of the small subunit ribosomal RNA gene sequence of *sarcocystis neurona* toward delineation of a definitive host and serologic studies.

M.S. Theses

(Unpublished)

Agricultural Economics

- Cao, Wuhong Cathy.* Cigarette consumption and the impact of health trend in East Asia.
- Huffman, Dana M.* An evaluation of the effects of the Community Reinvestment Act and other bank characteristics on levels of economic development in Kentucky.
- Widanski, Adam S.* An investigation into the export competitiveness of Polish fruits and vegetables: 1989-1994.

Agronomy

- McMurry, Stephen W.* Infiltration of fecal bacteria through intact soil profiles.
- Montgomery, Elizabeth D.* Herbicide and nitrate behavior in a shallow limestone aquifer.
- Ralston, Jennifer Lambert.* Movement of herbicides in shallow ground water in no-till double-cropped soybean.
- Saha, Hemedi M.* The influence of corn planting date, cover crop, and fertilizer N rate on the N nutrition of corn (*Zea mays L.*)
- Seither, Craig A.* Characterization of epoxy fatty acid synthesis in *Vernonia galamensis* and isolation of candidate cDNA clones.
- Yan, Xianghe.* Studies of signal sequences for cell-wall targeting in plants, and wall-targeting of a glutamine binding protein to effect cadmium tissue partitioning.

Animal Sciences

- Burchett, Stephen.* Effect of altering starch cook and resistant starch content of extruded food products on nutrient digestibility in the dog.

- Clarke, S.M.* Interaction between calf sex and calf genetic type for preweaning traits of calves varying in percentage Brahman breeding.
- Lu, Mei.* Influence of heat stress and calving on antioxidant activity in bovine blood.
- Reverte, Daniel.* Physiochemical and sensory properties of grass- and grain-fed beef treated with antioxidant and flavor agents.
- Silvio, Jennifer.* Influence of fiber fermentability on nutrient digestion in the dog.

Biosystems and Agricultural Engineering

- Tarhan, Sefa.* Evaluation of straw properties for biomass gasification.

Entomology

- Anderson, A.C.* Influence of soybean canopy closure on generalist predator abundance and predation of corn earworm eggs.
- Yu, Z.* Autodissemination of a baculovirus for suppression of beet armyworm, *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae).

Plant Pathology

- Smigell, Christopher.* Development of a polymerase chain reaction-based assay for the detection of *Peronospora tabacina* in tobacco transplants.

Veterinary Science

- McManus, Christina J.* Influence of glucose deprivation on the secretion of gonadotropins and prolactin in the mare.

Statewide Research

Research activities of the Kentucky Agricultural Experiment Station were conducted at Lexington, Princeton, Quicksand, and Owenton, and in counties throughout the state in 1996.

Efforts are constantly made to ensure that the research studies have application to the problems of all Kentucky farmers and other clientele groups. Locations of the experimental facilities provide conditions representative of most sections of the state.

Campus — Laboratories and specialized equipment for all research program areas.


Coldstream — Maine Chance — Spindletop Farms — Beef and dairy cattle, poultry, horses, sheep and swine, forages and grain crops, tobacco and turf.

South Farm — Fruits and vegetables, ornamentals.

UK Animal Research Center (Woodford County) — This farm was purchased in late 1991 as a location for development of state-of-the-art food animal research programs. The farm is in the initial phase of development as a research facility.

At Princeton (*Caldwell County*) the **Research and Education Center** facilities and the **West Kentucky Substation Farm** are devoted to research on grain crops, beef cattle, swine, fruits and vegetables, forages, and tobacco.

At Quicksand (*Breathitt County*) the **Robinson Substation** is the location of research on fruits and vegetables, or-



- 1 University of Kentucky, Lexington
- 2 Research and Education Center and West Kentucky Substation Farm, Princeton
- 3 Robinson Substation and Forest, Quicksand
- 4 Eden Shale Farm, Owenton

namentals, forages, grain crops, tobacco, and wood utilization. Quicksand is also the headquarters of **Robinson Forest**, which spreads over parts of Breathitt, Perry, and Knott counties and is the site of forestry and watershed management research.

The **Eden Shale Farm**, located in Owen County near Owenton, is where experimental and demonstration studies are conducted on forage crops, tobacco, and beef management.

Financial Statement

Statement of Current General Fund Income and Expenditures Fiscal Year 1996

Income

Federal Funds:	
Hatch Amended	\$ 3,700,984.00
Regional Research	776,071.00
McIntire-Stennis	392,434.00
Animal Health	80,701.00
Total Federal Funds	\$4,950,190.00
Non-Federal Funds	21,755,039.46
Total Funds	\$26,705,229.46

Expenditures

	<u>Federal</u>	<u>Non-Federal</u>	<u>Total</u>
Personal Services	\$4,176,223.74	\$13,838,894.07	\$18,015,117.81
Travel	106,963.34	103,856.72	210,820.06
Equipment	140,197.06	565,070.00	705,267.06
Other Operating Expenses	526,805.86	7,247,218.67	7,774,024.53
Total Expenditures	<u>\$4,950,190.00</u>	<u>\$21,755,039.46</u>	<u>\$26,705,229.46</u>

Staff

University of Kentucky Board of Trustees 1996

Edward T. Breathitt, Chairperson
Ted Bates

Kay Shropshire Bell
Paul W. Chellgren
Robert N. Clay
Merwin Grayson, Jr.
Jack Guthrie

James F. Hardymon
Robert P. Meriwether
Billy Joe Miles
Elissa M. Plattner
Steven S. Reed
C. Frank Shoop

Lois Combs Weinberg
Martin Welenken
Billy B. Wilcoxson

Faculty Members: Loys Mather,
Jimmy Jack Miller, & Deborah Powell
Student Member: Alan Aja

Agricultural Experiment Station

January 1 — December 31, 1996

Administration

Charles T. Wethington, Jr., President
Elisabeth Zinser, Chancellor
C.O. Little, Dean & Director
H.C. Owen, Treasurer

J.A. Boling, Associate Dean for Research & Associate Director

A.J. Hiatt, Associate Dean for Administration

J.D. Lawson, Assistant Director for Legal & Fiscal Affairs

W.O. Peterson, Director of Management Operations

Departments

Agricultural Communications

Craycraft, C.G., Director
Fehr, R.L., Professor
Weckman, R.D., Associate Professor
Witham, D.B., Associate Professor

Agricultural Economics

Robbins, L.W., Chairman & Professor
Bobst, B.W., Associate Professor
Debertin, D.L., Professor
Fleming, R., Assistant Professor
Freshwater, D., Associate Professor
Goetz, S.J., Associate Professor
Hall, H.H., Professor
Marchant, M.A., Associate Professor
Mather, L.L., Associate Professor
Pagoulatos, A., Professor
Reed, M.R., Professor
Skees, J.R., Professor

Agronomy

Smith, M.S., Chairman & Professor
Barnhisel, R.I., Professor
Barrett, M., Associate Professor
Blevins, R.L., Professor
Burton, H.R., Associate Professor
Bush, L.P., Professor
Calvert, J.R., Research Specialist
Chappell, J., Associate Professor
Chemiae, G.M., Professor
Collins, G.B., Professor
Collins, M., Professor
Cornelius, P.L., Professor
Coyne, M.S., Assistant Professor
Dinkins, R.D., Research Specialist
Dougherty, C.T., Professor
Egli, D.B., Professor
Evangelou, V.P., Professor
Evans, K.O., Research Specialist
Fanin, F.F., Research Associate
Grabau, L.J., Associate Professor
Grove, J.H., Associate Professor
Hildebrand, D.F., Associate Professor
Hunt, A.G., Associate Professor
Karathanasis, A., Professor
Kennedy, B.S., Research Specialist
Lauriault, L.M., Research Specialist
Legg, P.D., Professor
Nielsen, M.T., Professor
Pearce, R.C., Assistant Professor
Perfect, E., Assistant Professor
Pfeiffer, T.W., Professor
Phillips, A.D., Research Specialist
Phillips, T.D., Assistant Professor
Polge, N.D., Research Associate
Poneleit, C.G., Professor
Powell, A.J., Professor
Slack, C.H., Research Specialist
Taylor, N.L., Professor
TeKrony, D.M., Professor
Thom, W.O., Professor
Thomas, G.W., Professor
Thompson, J.A., Assistant Professor
Tomes, L.J., Research Specialist
Tutt, C.R., Research Specialist

Van Sanford, D.A., Professor
Wagner, G., Professor
Witt, W.W., Professor
Wood, J.M., Research Specialist

Animal Sciences

Kiracofe, G.H., Chairman & Professor
Aaron, D.K., Associate Professor
Barnes, T.B., Research Specialist
Boatright, W.L., Assistant Professor
Burris, R., Professor
Button, F.S., Jr., Research Specialist
Campbell, E.R., Research Specialist
Cantor, A.H., Associate Professor
Cromwell, G.L., Professor
Dawson, K.A., Professor
Deweese, W.P., Research Specialist
Edgerton, L.A., Associate Professor
Ely, D.G., Professor
Ford, M.J., Research Specialist
Harmon, D.L., Professor
Harmon, R.J., Professor
Hemken, R.W., Professor
Hicks, C.L., Professor
Hightshoe, B., Research Specialist
Jackson, J.A., Jr., Associate Professor
Langlois, B.E., Professor
Larson, B.T., Assistant Professor
Lawrence, L.M., Associate Professor
Lindemann, M.D., Associate Professor
Mitchell, G.E., Jr., Professor
Monegue, H.J., Research Specialist
Moody, W.G., Professor
O'Leary, J., Associate Professor
Parker, G.R., Professor
Randolph, J.H., Research Specialist
Schillo, K.K., Associate Professor
Silvia, W.J., Associate Professor
Straw, M.L., Research Specialist
Strobel, H.J., Associate Professor
Thrift, F.A., Professor
Tidwell, J., Assistant Professor*
Wang, C., Assistant Professor*
Xiong, Y., Associate Professor
Zavos, P.M., Professor

* Adjunct

Biosystems and Agricultural Engineering

Walton, L.R., Chairman & Professor
Bridges, T.C., Research Specialist
Colliver, D.G., Associate Professor
Day, G., Research Specialist
Duncan, G.A., Professor
Edwards, D.R., Associate Professor
Gates, R.S., Professor
McNeill, S.G., Research Specialist
Murphy, W.E., Professor
Nokes, S.E., Assistant Professor
Overhults, D.G., Professor
Payne, F.A., Professor
Priddy, K.T., Research Specialist
Ross, I.J., Professor
Shearer, S.A., Associate Professor
Swetnam, L.D., Research Specialist
Taraba, J.L., Professor

Turner, L.W., Professor
Warner, R.C., Associate Professor
Wells, L.G., Professor
Workman, S.R., Assistant Professor

Entomology

Pass, B.C., Chairman & Professor
Barney, R.J., Assistant Professor*
Brown, G.C., Associate Professor
Collins, J.T., Nursery Inspector
Dahlman, D.L., Professor
Haynes, K.F., Professor
Hunt, R., Assistant Professor*
Knapp, F.W., Professor
Moore, A.J., Associate Professor
Parr, J.C., Research Specialist
Potter, D.A., Professor
Sedlacek, J.D., Assistant Professor*
Webb, B.A., Assistant Professor
Webb, J.D., Research Specialist
Weston, P.A., Assistant Professor*
Wise, D.H., Professor
Yeagan, K.V., Professor

* Adjunct

Forestry

Muller, R.N., Chairman & Associate Professor
Arthur, M.A., Assistant Professor
Brown, D.L., Assistant Professor
Cremeans, D.W., Research Specialist
Kalisz, P.J., Associate Professor
Kimmerer, T.W., Associate Professor
Lacki, M.J., Associate Professor
Liu, C.J., Associate Professor
Paratley, R.J., Research Specialist
Pelkki, M.H., Assistant Professor
Ringe, J.M., Associate Professor
Stringer, J.W., Assistant Professor
Wagner, D.B., Associate Professor

Horticulture and Landscape Architecture

Ingram, D.L., Chairman & Professor
Anderson, R.G., Professor
Archbold, D.D., Associate Professor
Brown, G.R., Professor
Buxton, J.W., Associate Professor
Geneve, R.L., Associate Professor
Houtz, R.L., Associate Professor
Kemp, T.R., Professor
McNiel, R.E., Professor
Merkin, Z., Research Specialist
Nieman, T.J., Professor
Rowell, A.B., Assistant Professor
Snyder, J.C., Associate Professor
Weston, L.A., Associate Professor
Wolfe, D.E., Research Specialist

Nutrition and Food Science

Mercer, L.P., Chairman & Professor
Barth, M., Associate Professor
Chen, L.H., Professor
Chow, C.K., Professor
Glauert, H.P., Professor
Hennig, B., Professor

Plant Pathology

Smith, D.A., Chairman & Professor
Ammar, E.D., Research Specialist
Bachi, P.R., Research Specialist
Beale, J.W., Research Specialist
Ghabrial, S.A., Professor
Hendrix, J.W., Professor
Jarfors, U.E., Research Specialist
Maiti, I.B., Research Specialist
Nuckles, E.M., Research Specialist
Pirone, T.P., Professor
Sanger, M.P., Research Specialist
Schardl, C.L., Associate Professor
Shain, L., Professor
Shaw, J.G., Professor
Shepherd, R.J., Professor
Siegel, M.R., Professor
Thornbury, D.W., Biomedical Scientist
Vaillancourt, L.J., Assistant Professor
Wu, X., Research Specialist

Regulatory Services

Frye, W.W., Director & Professor
Beine, R.L., Laboratory Coordinator
Buckingham, D.T., Seed Regulatory Coordinator
Case, V.W., Soil Testing Coordinator
Coffey, D.S., Inspector
Cox, B.W., Inspector
Fabrizius, E.E., Seed Testing Coordinator
Flood, J.S., Inspector
Gibson, G.L., Inspector
Hatton, R., Milk Coordinator
Herald, F., Inspection Coordinator
Johnston, N.T., Inspector
Kirkland, D.L., Soil Testing Specialist
Lichtenberg, D., Lab Specialist
Mason, D.W., Inspector
Miller, E., Feed Coordinator

Prather, T.G., Inspector
Spencer, H.S., Auditor
Terry, D.L., Fertilizer Coordinator & Assistant Director
Thio, A.P., Lab Specialist
Thompson, C.D., Inspector
Troutman, D., Inspector

Robinson Substation

Morrison, M., Superintendent

Rural Sociology

Houglund, J.G., Chairman & Professor
Burmeister, L.L., Associate Professor
Coughenour, C.M., Professor
Dyk, P.A., Associate Professor
Garkovich, L.E., Professor
Greider, T.R., Associate Professor
Harris, R.P., Associate Professor
Maurer, R.C., Associate Chairman & Professor
Strang, N.L., Research Specialist
Swanson, L., Professor

Veterinary Science

Timoney, P.J., Chairman & Professor
Adams, M.H., Research Specialist
Allen, G.P., Professor
Artushin, S.C., Research Specialist
Bailey, E.F., Professor
Chambers, T.M., Associate Professor
Coogle, L.D., Research Specialist
Cook, R.F., Research Specialist
Cothran, E.G., Associate Professor
Davison, L.A., Research Specialist
Donahue, J.M., Professor
Dwyer, R.M., Assistant Professor
Fitzgerald, B.P., Associate Professor

Giles, R.C., Professor
Granstrom, D.E., Associate Professor
Graves, K.A., Assistant Professor
Hale, G., Librarian
Harkins, J.D., Research Specialist
Harrison, L.H., Professor
Henney, P.J., Research Specialist
Hong, C.B., Professor
Issel, C.J., Professor
Lai, A.C.-K., Research Specialist
Leach, R.B., Research Specialist
Lear, T.L., Research Specialist
Lehner, A.F., Research Specialist
Lyons, E.T., Professor
McCollum, W.H., Professor
McDowell, K.J., Associate Professor
Newman, D.H., Research Specialist
Petrites-Murphy, M.B., Associate Professor
Poonacha, K.B., Professor
Powell, D.G., Associate Professor
Smith, B.J., Research Specialist
Smith, R.A., Associate Professor
Swerczek, T.W., Professor
Timoney, J.F., Professor
Tobin, T., Professor
Tolliver, S.C., Research Specialist
Tramontin, R.R., Associate Professor
Vickers, M.L., Associate Professor
Williams, N.M., Assistant Professor
Woods, W.E., Research Specialist
Wulff-Strobel, C., Research Specialist
Yeargan, M.R., Research Specialist

West Kentucky Substation

Davis, D., Superintendent

Editor — Linda R. Kiesel
Department of Agricultural Communications

Designer — Dennis P. Duross
Department of Agricultural Communications

Mention of a trademark or proprietary product is for experimental purposes and does not constitute a guarantee or warranty by the Kentucky Agricultural Experiment Station, and does not imply its approval to the exclusion of other products that may also be suitable.

Look for the College of Agriculture on the World Wide Web at: <http://www.ca.uky.edu/>